

452

version

last name

first name

signature

McCord CH301
50070

Exam 2

Oct 10, 2017
Tuesday

Remember to refer to the Periodic Table handout that is separate from this exam copy.
There are many conversion factors and physical constants available there.

NOTE: Please keep this exam copy intact (all pages still stapled - including this cover page). You must turn in ALL the materials that were distributed. This means that you turn in your exam copy (name and signature included), bubble sheet, periodic table handout, and all scratch paper. Please also have your UT ID card ready to show as well.

This print-out should have 20 questions. Multiple-choice questions may continue on the next column or page – find all choices before answering.

001 6.0 points

What is the wavelength of a 4.50×10^{14} Hz light ray?

1. 0.288 nm
2. 666 nm
3. 882 nm
4. 0.441 nm
5. 456 nm
6. 0.664 nm
7. 0.882 nm
8. 992 nm

002 5.0 points

Which of the following best describes the effect of blue light on matter?

1. Electron excitation
2. Repulsion
3. No effect
4. Rotation
5. Ionization
6. Vibration

003 5.0 points

A particle confined to a one-dimensional box of length 480 nm for Ψ_3 has zero probability of being found:

1. Only 240 nm
2. 160 nm and 320 nm

3. 80 nm, 240 nm, and 160 nm

4. 160 nm and 240 nm

5. 80 nm, 240 nm, and 400 nm

004 3.0 points

Effective nuclear charge represents the attraction between the nucleus and the outer (valence) electrons.

1. False
2. True


005 5.0 points

Rank the following species by ionic radius, from largest to smallest: Cl^- , K^+ , Ca^{2+} , S^{2-} .

1. K^+ , Ca^{2+} , S^{2-} , Cl^-
2. S^{2-} , Cl^- , K^+ , Ca^{2+}
3. Ca^{2+} , K^+ , Cl^- , S^{2-}
4. Ca^{2+} , S^{2-} , Cl^- , K^+
5. S^{2-} , Cl^- , Ca^{2+} , K^+
6. K^+ , Cl^- , Ca^{2+} , S^{2-}

006 5.0 points

The second ionization of Fe is given by the reaction:



This electron is removed from the:

1. 3s subshell
2. 4p subshell
3. 3d subshell
4. 3p subshell
5. 4s subshell

6. 4d subshell

007 5.0 points

Which of the following atoms has an electronic structure that will weakly attract a magnetic field?

1. Ne
2. Li
3. Be
4. Zn

008 6.0 points

The following table shows the first six ionization energies for an unknown element:

#	I.E. (kJ/mol)
1	7.89×10^2
2	1.58×10^3
3	3.23×10^3
4	4.36×10^3
5	1.61×10^4
6	1.98×10^4

The unknown element is:

1. N
2. Si
3. B
4. Ne
5. O

009 6.0 points

Which of the following statement(s) is/are true regarding the Schrödinger equation and particle in a box?

- I. When $n = 2$ for a particle in a box, the particle has zero probability of being found in the center of the box
- II. There are an infinite numbers of solutions to the Schrödinger equation

III. The Schrödinger equation gives only positive solutions for the wavefunction (ψ)

IV. The angular momentum quantum number is one of the Schrödinger equation solutions for the hydrogen atom.

1. I and II
2. II only
3. I, III, and IV
4. I, II, and III
5. IV only
6. I and IV
7. I only
8. I, II, and IV

010 5.0 points

Which of the following is the best description for a $4p$ orbital of hydrogen?

1. A dumbbell shape with three spherical nodes and one angular node
2. Four concentric spheres with two angular nodes
3. Four concentric spheres with three radial nodes
4. A dumbbell shape with two spherical nodes and one angular node
5. A dumbbell shape with one spherical node and three angular nodes
6. Four concentric spheres with one angular node
7. A dumbbell shape with four spherical nodes and one angular node

8. A dumbbell shape with one spherical node and two angular nodes

011 5.0 points

Which of the following statements is correct?

1. Ionization energy consistently increases as you approach the bottom left of the periodic table

2. Ionization energy consistently increases as you approach the top right of the periodic table

3. Phosphorus has a higher ionization energy than Sulfur because there are more electrons in the p subshell in Phosphorus

4. Magnesium has a higher ionization energy than Aluminum because the full s subshell of Magnesium is more stable than the partially filled p subshell of Aluminum

5. Sulfur has a higher ionization energy than Phosphorus because there are more electrons in the p subshell in Sulfur

6. None of the above

012 3.0 points

Which of the following is the correct name for the chemical formula: $(\text{NH}_4)_2\text{SO}_4$?

1. Ammonium sulfite

2. Nitrohydrogen sulfurous oxide

3. Nitrohydrogen sulfuric oxide

4. Ammonium sulfate

5. Ammonium sulfurous oxide

013 3.0 points

Which orbital corresponds to the following quantum number set:

$$m_s = -1/2, \quad m_l = -2, \quad n = 6, \quad l = 3$$

1. 6f

2. 6d

3. 6p

4. 6f and 6d are both possible

5. All choices are possible

6. 3s

014 5.0 points

What is the correct shorthand electron configuration for copper?

1. $[\text{Kr}]4d^45s^2$

2. $[\text{Ar}]3d^94s^2$

3. $[\text{Ar}]3d^44s^2$

4. $[\text{Ar}]3d^{10}4s^1$

5. $[\text{Kr}]4d^55s^1$

6. $[\text{Ar}]3d^54s^1$

015 6.0 points

What is the change in energy for an excited hydrogen electron falling from the $n = 4$ to the $n = 2$ energy level? Is light absorbed or emitted through this process?

1. 4.09×10^{-17} J, absorbed

2. -4.09×10^{-17} J, emitted

3. -2.06×10^{-17} J, absorbed

4. -2.06×10^6 J, absorbed

5. -4.09×10^{-19} J, emitted

6. 2.06×10^{-17} J, absorbed

016 5.0 points

Which set of quantum numbers is possible for an electron in the p subshell found in the

ground state of As?

1. $n = 3, \ell = 2, m_\ell = +2, m_s = +\frac{1}{2}$
2. $n = 4, \ell = 1, m_\ell = +1, m_s = -\frac{1}{2}$
3. $n = 1, \ell = 1, m_\ell = +2, m_s = -\frac{1}{2}$
4. $n = 4, \ell = 2, m_\ell = +1, m_s = -\frac{1}{2}$
5. $n = 4, \ell = 0, m_\ell = 0, m_s = +\frac{1}{2}$
6. $n = 4, \ell = 4, m_\ell = +2, m_s = +\frac{1}{2}$

017 5.0 points

Consider the following two gaseous samples at STP:

Sample A: 10L Hydrogen

Sample B: 3L Hydrogen

Each sample is excited with electricity and the light emitted is refracted through a dispersive prism. Which of the following best describes the difference between the two samples?

1. The emission spectrum for Sample A will show continuous lines in lower wavelengths
2. Both emission spectra will be identical
3. The emission spectrum for Sample A will show more discrete lines in higher wavelengths
4. The emission spectrum for Sample A will be brighter than Sample B
5. The emission spectrum for Sample A will show more discrete lines in lower wavelengths

018 6.0 points

Which of the following best matches the quantum mechanical evidence to its conclusion?

1. Particle in a box: an electron can be found at any distance between 0 and L, with the highest probability being at L/n
2. Rydberg Energy Levels: the energy difference between consecutive n values decreases

linearly as n increases

3. Absorption spectrum: the line spectra of any gas will show the same characteristic wavelengths

4. Emission spectrum: an excited sample of hydrogen gas releases characteristic wavelengths of light when its electrons fall from a high discrete energy level to a low discrete energy level

019 5.0 points

Which of the following is the correct shorthand electron configuration for Sb?

1. $[\text{Ar}]4s^23d^{10}4p^2$
2. $[\text{Kr}]5s^24d^{10}5p^1$
3. $[\text{Kr}]5s^25p^1$
4. $[\text{Kr}]5s^25p^2$
5. $[\text{Ar}]4s^24d^{10}4p^1$
6. $[\text{Kr}]5s^24d^{10}5p^3$
7. $[\text{Ar}]4s^23d^{10}4p^1$
8. $[\text{Kr}]5s^24d^{10}5p^2$

020 6.0 points

A 37.0 nm beam of light is shined on a chromium surface. What is the maximum kinetic energy of the excited electrons? The work function of chromium is 4.50 eV.

1. No electrons are emitted
2. 1×10^{-17} J
3. 2.08×10^{-17} J
4. -7.16×10^{-19} J
5. 5.37×10^{-12} J
6. 4.65×10^{-18} J

7. -4.50 J

8. 5.37×10^{-15} J