HW13 - First Law & Calorimetry **Question 1** 1 pts A 100 W electric heater (1 W = 1 J/s) operates for 11 min to heat the gas in a cylinder. At the same time, the gas expands from 1 L to 6 L against a constant atmospheric pressure of 3.527 atm. What is the change in internal energy of the gas? 67.79 kJ 62.47 kJ 48.37 kJ O 64.21 kJ **Question 2** 1 pts The definition of internal energy is $\Delta U = q + w$ Which of these three values are state functions? Select all of the correct answers. W \Box ΔU □ q **Question 3** 1 pts When 2.00 kJ of energy is transferred as heat to nitrogen in a cylinder fitted with a piston with an external pressure of 2.00 atm, the nitrogen gas expands from 2.00 to 5.00 L. What is the change in internal energy of this system? +2.61 kJ +1.39 kJ -0.608 kJ \bigcirc 0 -2.61 kJ **Question 4** 1 pts A system had 150 kJ of work done on it and its internal energy increased by 60 kJ. How much energy did the system gain or lose as heat? The system gained 60 kJ of energy as heat. The system lost 210 kJ of energy as heat. The system gained 90 kJ of energy as heat. The system lost 90 kJ of energy as heat. The system gained 210 kJ of energy as heat. **Question 5** 1 pts If a process is carried out at constant pressure and the volume of the system decreases, then ΔV is [Select] and the work is [Select] **Question 6** 1 pts Which of the following will best help determine the direction of heat flow in a system? internal energy pressure work temperature enthalpy **Question 7** 1 pts Which of the following statements concerning the first law of thermodynamics is/are true? Select all of the correct answers. The universe is an isolated system. The internal energy of the universe is always increasing. Internal energy lost by a system is always gained by the surroundings. **Question 8** 1 pts What is the value of work when a piston of volume 0.2 L expands against an external pressure of 200 kPa to a volume of 3.4 L? 3.40 kJ 3.40 kJ -640 J 640 J **Question 9** 1 pts When 4.00 kJ of energy is transferred as heat to nitrogen in a cylinder fitted with a piston at an external pressure of 3.00 atm, the nitrogen gas expands from 1.00 L to 4.00 L against this constant pressure. What is ΔU for the process? Note: 1 L*atm = 0.1013 kJ. +4.91 kJ -0.912 kJ +3.09 kJ -4.91 kJ **Question 10** 1 pts A piece of metal with a mass of 22 g at 92 °C is placed in a calorimeter containing 53.7 g of water at 21 °C. The final temperature of the mixture is 55.3 °C. What is the specific heat capacity of the metal? Assume that there is no energy lost to the surroundings. O -9.5 J g⁻¹ °C⁻¹ \bigcirc -1.3 x 10⁴ J g⁻¹ °C⁻¹ ○ 9.5 J g⁻¹ °C⁻¹ \bigcirc 1.3 x 10⁴ J g⁻¹ °C⁻¹ **Question 11** 1 pts Consider the following specific heat capacities: $H_2O(s) = 2.09 \text{ J/g} \cdot ^{\circ}C$ $H_2O(I) = 4.18 \text{ J/g} \cdot ^{\circ}C$ $H_2O(g) = 2.03 \text{ J/g} \cdot ^{\circ}C$ The heat of fusion for water is 334 J/g and its heat of vaporization is 2260 J/g. Calculate the amount of heat required to convert 93 g of ice at -36°C completely to liquid water at 35°C. 52 kJ 21 kJ 38 kJ 7 kJ **Question 12** 1 pts The specific heat for liquid argon and gaseous argon is 25.0 J/mol·°C and 20.8 J/mol·°C, respectively. The enthalpy of vaporization of argon is 6506 J/mol. How much energy is required to convert 1 mole of liquid Ar from 5°C below its boiling point to 1 mole of gaseous Ar at 5°C above its boiling point? 6631 J 6735 J O 229 J 125 J 6610 J **Question 13** 1 pts Carbon monoxide reacts with oxygen to form carbon dioxide by the following reaction: $2CO(g) + O_2(g) \rightarrow 2CO_2(g)$ ΔH for this reaction is -135.28 kcal. How much heat would be released if 12.0 moles of carbon monoxide reacted with sufficient oxygen to produce carbon dioxide? Use only the information provided in this question. 1620 kcal 812 kcal 135 kcal 412 kcal **Question 14** 1 pts What mass of liquid ethanol (C₂H₅OH) must be burned to supply 500 kJ of heat? The standard enthalpy of combustion of ethanol at 298 K is -1368 kJ/mol. 16.8 g 10.9 g 29.7 g 126 g **Question 15** 1 pts Burning 1 mol of methane in oxygen to form CO₂ (g) and H₂O (g) produces 803 kJ of energy. How much energy is produced when 3 mol of methane is burned? 1606 kJ 2409 kJ 803 kJ 268 kJ **Question 16** 1 pts Consider the following chemical equation: $C_3H_8(g) + 5O_2(g) \rightarrow 3CO_2(g) + 4H_2O(I)$ $\Delta H = -2220 \text{ kJ/mol rxn}$ How much thermal energy is given off when 11.0 g of propane gas (C₃H₈) is burned at constant pressure? 2220 kJ 555 kJ 1670 kJ 26.0 kJ **Question 17** 1 pts For a certain reaction at constant pressure, the change in internal energy is -52 kJ. In addition, the system does 46 kJ of expansion work. What is ΔH for this process? -98 kJ 98 kJ -6 kJ 6 kJ **Question 18** 1 pts If the products of a reaction have higher energy than the reactants, then the reaction... is exothermic. is endothermic. must be spontaneous. is not spontaneous. **Question 19** 1 pts The specific heats and densities of several materials are given below: Specific Heat (cal/g·°C) Density (g/cm³) Material **Brick** 0.220 2.0 0.270 2.7 Concrete 7 Steel 0.118 Water 1.00 1.00 Calculate the change in temperature produced by the addition of 1 kcal of heat to 100 g of steel. 37.0°C 84.7°C 1.18°C 1.43°C **Question 20** 1 pts A 1.00 g sample of n-hexane (C_6H_{14}) undergoes complete combustion with excess O_2 in a bomb calorimeter. The temperature of the 1502 g of water surrounding the bomb rises from 22.64°C to 29.30°C. The heat capacity of the hardware component of the calorimeter (everything that is not water) is 4042 J/°C. What is ΔU for the combustion of $n-C_6H_{14}$? One mole of $n-C_6H_{14}$ is 86.1 g. The specific heat of water is 4.184 J/g·°C. -1.15 x 10⁴ kJ/mol \circ -9.96 x 10³ kJ/mol -4.52 x 10³ kJ/mol -5.92 x 10³ kJ/mol **Question 21** 1 pts When 0.485 g of compound X is burned completely in a bomb calorimeter containing 3000 g of water, a temperature rise of 0.285°C is observed. What is ΔU of the reaction for the combustion of compound X? The hardware component of the calorimeter has a heat capacity of 3.81 kJ/°C. The specific heat of water is 4.184 J/g·°C, and the MW of X is 56.0 g/mol. 538 kJ/mol -4660 kJ/mol -538 kJ/mol 4660 kJ/mol **Question 22** 1 pts Nitric acid can be manufactured in a multi-step process, during which nitric oxide is oxidized to create nitrogen dioxide. $2NO(g) + O_2(g) \rightarrow 2NO_2(g)$ Calculate the standard reaction enthalpy for the above reaction using the following thermodynamic data. $N_2(g) + O_2(g) \rightarrow 2NO(g)$ $\Delta H^{\circ}_1 = 180.5 \text{ kJ/molrxn}$ $N_2(g) + 2O_2(g) \rightarrow 2NO_2(g)$ $\Delta H_2^\circ = 66.4 \text{ kJ/molrxn}$ -100.3 kJ/mol rxn -252.4 kJ/mol rxn -246.9 kJ/mol rxn -114.1 kJ/mol rxn Question 23 1 pts Calculate the standard reaction enthalpy for the following chemical equation. $CH_4(g) + H_2O(g) \rightarrow CO(g) + 3H_2(g)$ Use the following thermochemical equations to solve for the change in enthalpy. $2H_2\left(g\right) + CO\left(g\right) \rightarrow CH_3OH\left(I\right)$ $\Delta H^{\circ} = -128.3 \text{ kJ/mol}$ $2CH_{4}\left(g\right)+O_{2}\left(g\right)\rightarrow2CH_{3}OH\left(I\right) \hspace{1.5cm}\Delta H^{\circ}=-328.1\text{ kJ/mol}$ $2H_2(g) + O_2(g) \rightarrow 2H_2O(g)$ $\Delta H^{\circ} = -483.6 \text{ kJ/mol}$ +206.1 kJ/mol +42.0 kJ/mol +216 kJ/mol +155.5 kJ/mol **Question 24** 1 pts Calculate the standard enthalpy change for the following chemical equation. $2HCI(g) + F_2(g) \rightarrow 2HF(I) + CI_2(g)$ Use the following thermochemical equations to solve for the change in enthalpy. $4HCI(g) + O_2(g) \rightarrow 2H_2O(I) + 2CI_2(g)$ $\Delta H^{\circ} = -202.4 \text{ kJ/mol rxn}$ $\Delta H^{\circ} = -600.0 \text{ kJ/mol rxn}$ $\frac{1}{2}$ H₂(g) + $\frac{1}{2}$ F₂(g) \rightarrow HF (I) $H_2(g) + \frac{1}{2}O_2(g) \rightarrow H_2O(I)$ ΔH° = -285.8 kJ/mol rxn -1088.2 kJ/mol rxn +516.6 kJ/mol rxn -1015.4 kJ/mol rxn -516.6 kJ/mol rxn -1587.2 kJ/mol rxn -1116.6 kJ/mol rxn +1116.6 kJ/mol rxn +1587.2 kJ/mol rxn +1015.4 kJ/mol rxn +1088.2 kJ/mol rxn **Question 25** 1 pts Calculate the standard enthalpy change for the following chemical equation. $4FeO (s) + O_2 (g) \rightarrow 2Fe_2O_3 (s)$ Use the following thermochemical equations to solve for the change in enthalpy. Fe (s) + $\frac{1}{2}$ O₂ (g) \rightarrow FeO (s) $\Delta H = -269 \text{ kJ/mol}$ 2Fe (s) + 3/2 O_2 (g) \rightarrow Fe₂ O_3 (s) $\Delta H = -825 \text{ kJ/mol}$ 574 kJ/mol -556 kJ/mol -574 kJ/mol -2726 kJ/mol 556 kJ/mol **Question 26** 1 pts Calculate the enthalpy change for the following chemical equation. $2SO_2(g) + O_2(g) \rightarrow 2SO_3(g)$ Use the following thermochemical data to solve for the change in enthalpy. ΔH_f for $SO_2(g) = -16.9 \text{ kJ/mol}$ ΔH_f for $SO_3(g) = -21.9 \text{ kJ/mol}$ -10.0 kJ/mol rxn +5.0 kJ/mol rxn -5.0 kJ/mol rxn -77.6 kJ/mol rxn **Question 27** 1 pts Which of the following substances have ΔH_f° = 0? Select all of the correct answers. C (s, graphite) ☐ HCl (aq) C (s, diamond) \Box $F_2(g)$ ☐ HCl (g) □ Na (s) **Question 28** 1 pts Calculate the average S–F bond energy in SF₆ using the following ΔH_f values: $SF_6(g) = -1209 \text{ kJ/mol}$ S(g) = 279 kJ/molF(g) = 79 kJ/mol327 kJ/mol bonds 289 kJ/mol bonds 582 kJ/mol bonds 196 kJ/mol bonds 416 kJ/mol bonds **Question 29** 1 pts Using the bond energy data provided, calculate ΔH for the following reaction: $H_2(g) + Cl_2(g) \rightarrow 2HCl(g)$ Bond Bond Energy (kJ/mol) H-H436 CI-CI 242 H-CI 432 -246 kJ/mol 186 kJ/mol 246 kJ/mol -186 kJ/mol **Question 30** 1 pts The standard molar enthalpy of formation of NH₃ (g) is -46.11 kJ/mol. What is the standard molar internal energy of formation of NH₃ (g)? -43.63 kJ/mol -2525 kJ/mol -48.59 kJ/mol 2433 kJ/mol