

HW03 - Chemical Equilibria

⚠ This is a preview of the draft version of the quiz


Started: Oct 1 at 12:06pm

Quiz Instructions

Question 1

1.2 pts

When the chemical reaction



is at equilibrium, which of the following is true?

- the sum of the concentrations of A and B equals the sum of the concentrations of C and D
- all four concentrations are equal
- both the forward and reverse reactions have stopped
- neither the forward nor the reverse reactions have stopped

Question 2

1.2 pts

Explain why equilibrium constants are dimensionless.

- Activities (which are dimensionless) are actually what should be used in the mass action expression and therefore equilibrium constants. Concentration and pressure *values* are used in place of activities of species. Therefore true equilibrium constants have no units.
- This is a trick question. Equilibrium constants have units that involve some multiple of atmospheres or moles per liter.
- They are dimensionless because the pressures or concentrations we put in are all for the substances in their standard states.


- They are not really dimensionless, but we must treat them as such in order to be able to take $\ln(K)$ in the expression:

$$\Delta G^\circ = -RT \ln K$$

Question 3

1.2 pts

The expression for K_c for the reaction



at equilibrium is:

$\frac{[\text{NO}][\text{H}_2\text{O}]}{[\text{NH}_3][\text{O}_2]}$

$[\text{NH}_3]^4[\text{O}_2]^5$




$\frac{[\text{NO}]^4[\text{H}_2\text{O}]^6}{[\text{NH}_3]^4[\text{O}_2]^5}$

$\frac{[\text{NH}_3]^4[\text{O}_2]^5}{[\text{NO}]^4[\text{H}_2\text{O}]^6}$

Question 4

1.2 pts

Consider the following reactions at 25°C:



Which compound is most likely to dissociate and give $\text{O}_2(\text{g})$ at 25°C?

H_2O

NO


CO

CO₂


Question 5

1.2 pts

At 600°C, the equilibrium constant for the reaction



is 2.8. Calculate the equilibrium constant for the reaction



1.7

1.1


0.36

0.60

Question 6

1.2 pts

Consider the reaction



What is the form of the equilibrium constant K_c for this reaction?

$\frac{[\text{Hg}]^2[\text{O}_2]}{[\text{HgO}]^2}$

$[\text{Hg}]^2 [\text{O}_2]$


$\frac{[\text{O}_2]}{[\text{HgO}]^2}$

[O₂]

Question 7

1.2 pts

$K_c = 2.6 \times 10^8$ at 825 K for the reaction




The equilibrium concentration of H₂ is 0.0020 M and S₂ is 0.0010 M. What is the equilibrium concentration of H₂S?

- 0.10 M
- 0.0010 M
- 10 M
- 1.0 M

Question 8

1.2 pts

Consider the reaction below




At 1000 K the equilibrium pressures of the three gases in one mixture were found to be 0.562 atm SO₂, 0.101 atm O₂, and 0.332 atm SO₃. Calculate the value of K_p for the reaction.

- 0.171
- 2.64
- 3.46
- 0.289

Question 9

1.2 pts

Consider the following reaction:



Calculate K_c for this reaction at 100°C .

 19.7 0.0784 7440 73.5**Question 10**


1.2 pts

Calculate the equilibrium constant at 25°C for a reaction for which $\Delta G^\circ = -4.22 \text{ kcal/mol}$.

 -1240.51 1240.51 2481.02 620.254**Question 11**

1.2 pts

The reaction



has an equilibrium constant of 3.7×10^{-3} . Consider a reaction mixture with:

$$[A] = 2.0 \times 10^{-2} \text{ M}$$

$$[B] = 1.7 \times 10^{-4} \text{ M}$$

$$[C] = 2.4 \times 10^{-6} \text{ M}$$

$$[D] = 3.5 \times 10^{-3} \text{ M}$$


Which of the following statements is definitely true?

- The system is at equilibrium.
- The forward reaction will occur to a greater extent than the reverse reaction until equilibrium is established.
- The reverse reaction will occur to a greater extent than the forward reaction until equilibrium is established.
- No conclusions about the system can be made without additional information.

Question 12

1.2 pts

The reaction




has an equilibrium constant (K_c) of 4.0×10^8 at 25°C . What will eventually happen if 44.0 moles of NH_3 , 0.452 moles of N_2 , and 0.108 moles of H_2 are put in a 10.0 L container at 25°

- Nothing. The system is at equilibrium.
- More N_2 and H_2 will be formed.
- It is impossible to know what will happen unless we know what the equilibrium constant is at 298 K.
- More NH_3 will be formed.

Question 13

1.2 pts

Consider the reaction:



If the initial concentration of $\text{Ni}(\text{CO})_4(\text{g})$ is 1.0 M and x is the equilibrium concentration of $\text{CO}(\text{g})$, what is the correct equilibrium relation?

$K_c = \frac{x^4}{\left(1.0 - \frac{x}{4}\right)}$

$K_c = \frac{4x}{(1.0 - 4x)}$


$K_c = \frac{256x^4}{(1.0 - 4x)}$

$K_c = \frac{x^5}{\left(1.0 - \frac{x}{4}\right)}$

Question 14

1.2 pts

Suppose the reaction



has an equilibrium constant $K_c = 49$ and the initial concentrations of H_2 and I_2 is 0.5 M and of HI is 0.0M. Which of the following is the correct value for the final concentration of $\text{HI}(\text{g})$?

0.778 M

0.599 M


0.250 M

0.219 M

Question 15

1.2 pts

The system




is at equilibrium at a fixed temperature with a partial pressure of H_2 of 0.200 atm, a partial pressure of I_2 of 0.200 atm, and a partial pressure of HI of 0.100 atm. An additional 0.26 atm pressure of HI is admitted to the container, and it is allowed to come to equilibrium again. What is the new partial pressure of HI?

- 0.152 atm
- 0.104 atm
- 0.464 atm
- 0.360 atm

Question 16

1.2 pts

At 990°C , $K_c = 1.6$ for the reaction




How many moles of $\text{H}_2\text{O}(\text{g})$ are present in an equilibrium mixture resulting from the addition of 1.00 mole of H_2 , 2.00 moles of CO_2 , 0.75 moles of H_2O , and 1.00 mole of CO to a 5.00 liter container at 990°C ?

- 0.60 mol
- 1.1 mol
- 1.7 mol
- 1.0 mol

Question 17

1.2 pts

What happens to the concentration of NO(g) when the total pressure on the reaction below is increased (by compression) when it is at equilibrium?




- it remains the same
- it is impossible to tell
- it increases
- it decreases

Question 18

1.2 pts

Consider the following reaction:




where $\Delta H_{rxn} = -198$ kJ. The amount of SO₂(g) at equilibrium increases when...

- SO₃ is removed.
- the volume is increased.
- the temperature is decreased.
- more oxygen is added.

Question 19

1.2 pts

Suppose the reaction mixture




is at equilibrium at a given temperature and pressure. The pressure is then increased at constant temperature by compressing the reaction mixture, and the mixture is then allowed to reestablish equilibrium. At the new equilibrium...

- the nitrogen is used up completely.
- there is less ammonia present than there was originally.
- there is the same amount of ammonia present as there was originally.
- there is more ammonia present than there was originally.

Question 20

1.2 pts

Consider the system:




at equilibrium at 25°C. If this is an exothermic reaction and the temperature was raised, would the equilibrium be shifted to produce more N_2O_5 or more N_2O_4 ?

- there would be no change
- more N_2O_5
- it is impossible to tell
- more N_2O_4

Question 21

1.2 pts

The system




is at equilibrium at some temperature. At equilibrium, a 4.00L vessel contains 1.00 mole CO_2 , 1.00 mole H_2 , 2.40 moles H_2O , and 2.40 moles CO . How many moles of CO_2 must be added to this system to bring the equilibrium CO concentration to 0.669 mol/L?

- 0.993 moles
- 0.429 moles
- 0.069 moles
- 0.498 moles

Question 22

1.2 pts

The figure below represents a reaction at 298 K.



Based on the figure, which of the following statements (if any) are FALSE?

- None of the other statements are false.
- At point C, the system is at equilibrium.

For this reaction, ΔG° is negative.


At point D, the reaction will move toward the reactants to get to equilibrium.

At point B, $Q < K$.

Question 23

1.2 pts

Given the hypothetical reaction:



Predict what will happen when 1.0 mol Y is placed into an evacuated container.

ΔG° will decrease until $\Delta G^\circ = 0$.

Q will increase until $Q = K$.


Nothing. The products are already formed, so no reaction occurs.

Q will decrease until $Q = K$.

Question 24

1.2 pts

Consider the reaction:



Which of the following is a possible value of K for this reaction?

10^{70}

0.56

-0.56

10^{-70}

Question 25

1.2 pts

The equilibrium constant K for the synthesis of ammonia is 6.8×10^5 at 298 K. What will K be for the reaction at 375 K?



- 326
- 6.85×10^5
- 6.75×10^5
- 1.42×10^9

Not saved

Submit Quiz