

HW07 - Kinetics

ⓘ This is a preview of the published version of the quiz

Started: Nov 8 at 5:49pm

Quiz Instructions

Question 1

0.5 pts

Consider the reaction:




What is the overall order of the reaction and the order with respect to $[\text{O}_3]$?

- 3 and 2
- 1 and 2
- 2 and 2
- 1 and 3

Question 2

0.5 pts

When the reaction below:




is proceeding under conditions such that 0.015 mol/L of N_2O is being formed each second, the rate of the overall reaction is _____ and the rate of change for NO is _____.

- 0.030 M/s; -0.005 M/s
- 0.015 M/s; -0.045 M/s
- 0.015 M/s; -0.005 M/s
- 0.015 M/s; 0.045 M/s

Question 3

1 pts

What is the rate law for the reaction below:



if the following data were collected?

Exp	$[\text{A}]_0$	$[\text{B}]_0$	$[\text{C}]_0$	Initial Rate
1	0.4	1.2	0.7	2.32×10^{-3}
2	1.3	1.2	0.9	7.54×10^{-3}
3	0.4	4.1	0.8	9.25×10^{-2}
4	1.3	1.2	0.2	7.54×10^{-3}

rate = $1.79 \times 10^{-3} [B]^2 [C]$

rate = $5.37 \times 10^{-3} [A] [B]^3$

rate = $1.49 \times 10^{-3} [B]^3 [C]$


rate = $4.48 \times 10^{-3} [A] [B]^2 [C]$

rate = $3.36 \times 10^{-3} [A] [B]^3$

Question 4

1 pts

A chemical reaction is expressed by the balanced chemical equation:



Consider the data below:

exp	[A] ₀	[B] ₀	initial rate (M/min)
1	0.15	0.15	0.00110363
2	0.15	0.3	0.0044145
3	0.3	0.3	0.008829

Find the rate law for the reaction.

rate = $k [B]^2$

rate = $k [A] [B]$

rate = $k [A]^2 [B]$

rate = $k [A] [B]^2$

Question 5

0.5 pts

Calculate the value of the rate constant (k) for the reaction in question 4.

0.000166

0.00736

0.00110

0.327

Question 6

0.5 pts

If the initial concentrations of both A and B are 0.31 M for the reaction in questions 4 and 5, at what initial rate is C formed?

0.00974 M/min

0.101 M/min


-0.00974 M/min

0.0314 M/min

Question 7

1 pts

We know that the rate expression for the reaction below:



at a certain temperature is $\text{rate} = [\text{NO}]^2 [\text{O}_2]$. We carry out two experiments involving this reaction at the same temperature, but in the second experiment the initial concentration of NO is doubled while the initial concentration of O_2 is halved. The initial rate in the second experiment will be how many times that of the first?

2

1

8

4

Question 8

1 pts

Consider the data collected for a chemical reaction between compounds A and B that is first order in A and first order in B:

rxn	[A] ₀	[B] ₀	rate (M/s)
1	0.2	0.05	0.1
2	?	0.05	0.4
3	0.4	?	0.8

From the information above for 3 experiments, determine the missing concentrations of A and B. Answers should be in the order [A] then [B].

0.40 M; 0.10 M

0.80 M; 0.20 M

0.20 M; 0.80 M

0.80 M; 0.10 M

0.40 M; 0.20 M

Question 9

0.5 pts

For a reaction that is zero-order overall...

the rate constant is zero.

the reactant concentration does not change with time.


the activation energy is zero.

- the rate does not change during the reaction.

Question 10

0.5 pts

Consider the reaction below:




If it is 1st order in A and 0th order in B, a plot of $\ln[A]$ vs time will have a slope that is...

- decreasing exponentially.
- increasing exponentially.
- constant.
- slowly increasing.

Question 11

1 pts

Consider the reaction below:




If it has a half-life of 1.6 sec, how long will it take a system with $[\text{H}_2\text{CO}_3]_0$ of 2M to reach $[\text{H}_2\text{CO}_3]$ of 125mM?

- 6.4 sec
- Not enough information is given.
- 3.2 sec
- 2.9 sec

Question 12

1 pts

At a certain fixed temperature, the reaction below:




is found to be first order in the concentration of A and zeroth order in the concentration of B. The reaction rate constant is 0.05s^{-1} . If 2.00 moles of A and 4.00 moles of B are placed in a 1.00 liter container, how many seconds will elapse before the concentration of A has fallen to 0.30 moles/liter?

- 37.94 sec
- 2.83 sec
- There is not enough information to answer.
- 10.22 sec

Question 13

0.5 pts

The reaction below:




is observed to obey first-order kinetics. Which of the following plots should give a straight line?

- [A] vs t
- [A] vs t^{-1}
- $\ln[A]$ vs k^{-1}
- $\ln[A]$ vs k
- $\ln[A]$ vs t^{-1}
- [A] vs k
- $\ln[A]$ vs t

Question 14

1 pts

For the reaction below:



at 800K, a plot of $\ln[\text{cyclobutane}]$ vs t gives a straight line with a slope of -1.6 s^{-1} . Calculate the time needed for the concentration of cyclobutane to fall to 1/16 of its initial value.

- 1.6 sec
- 1.3 sec
- 1.7 sec
- 0.63 sec

Question 15

1 pts

The initial concentration of the reactant A in a first-order reaction is 1.2 M. After 69.3 sec, the concentration has fallen to 0.3 M. What is the rate constant k?

- 0.01 s^{-1}
- 0.02 s^{-1}
- 0.2 s^{-1}
- not enough information

Question 16

1 pts




A reaction is found to be first order with respect to one of the reactant species, A. When might a plot of $\ln[A]$ vs time NOT yield a straight line?

- if the reaction has any significant backward rate
- All of the other answers could be correct.
- if the reaction comes to equilibrium
- when the rate also depends on the concentration of another reactant as well

Question 17

0.5 pts

Consider the following elementary reactions:




Identify the molecularity of each reaction respectively.

- all three elementary reactions are bimolecular
- bimolecular, unimolecular, termolecular
- it is impossible to know without knowing the overall reaction for each
- tetramolecular, termolecular, pentamolecular

Question 18

1 pts

A and B react to form C according to the single step reaction below:





Which of the following is the correct rate equation for [B] and the correct units for the rate constant of this reaction?

- $\frac{\Delta[\text{B}]}{\Delta t} = -\frac{2k[\text{A}][\text{B}]}{[\text{C}]}; \frac{1}{\text{M}\cdot\text{s}}$
- $\frac{\Delta[\text{B}]}{\Delta t} = -2k[\text{A}][\text{B}]; \frac{1}{\text{M}\cdot\text{s}}$
- $\frac{\Delta[\text{B}]}{\Delta t} = -2k[\text{A}][\text{B}]^2; \frac{1}{\text{M}^2\cdot\text{s}}$
- $\frac{\Delta[\text{B}]}{\Delta t} = -k[\text{A}][\text{B}]^2; \frac{1}{\text{M}^2}$

Question 19

1 pts

Consider the mechanism below:



What is the rate law?

-

rate = $k_1[\text{NO}_2][\text{F}_2]$

rate = $k_1k_2[\text{NO}_2]^2$

rate = $k_2[\text{NO}_2]^2$



rate = $k_1[\text{NO}_2\text{F}][\text{F}_2]$

rate = $k_2[\text{NO}_2][\text{F}]$

Question 20

1 pts

Determine the overall balanced equation for a reaction having the following proposed mechanism:



and write an acceptable rate law.

$2\text{B}_2 \longrightarrow \text{E}_3 + \text{D}$; rate = $k[\text{B}_2]^2$

$\text{A} + \text{B}_2 \longrightarrow \text{C}_2 + \text{D}$; rate = $k[\text{A}][\text{B}_2]$


$\text{A} + \text{B}_2 \longrightarrow \text{C}_2 + \text{D}$; rate = $k[\text{B}_2]^2$

$\text{E}_3 + \text{A} \longrightarrow \text{B}_2 + \text{C}_2$; rate = $k[\text{E}_3][\text{A}]$



Question 21

1 pts

Consider the reaction below:



The proposed mechanism of this reaction is:



What is the rate of the overall reaction?

rate = $k_1k_2[\text{I}_2][\text{H}_2]$

rate = $k_2[\text{I}]^2[\text{H}_2]$

rate = $\frac{k_1k_2}{k_{-1}}[\text{I}_2][\text{H}_2]$

rate = $\frac{k_{-1}k_2}{k_1}[\text{I}_2][\text{H}_2]$

rate = $\frac{k_1k_2}{k_{-1}}[\text{I}]^2[\text{H}_2]$

Question 22

1 pts

A reaction rate increases by a factor of 655 in the presence of a catalyst at 37°C. The activation energy of the original pathway is 106 kJ/mol. What is the activation energy of the new pathway, all other factors being equal?

- 89.3 J/mol
- 16,600 J/mol
- 89.3 kJ/mol
- 16,600 kJ/mol


Question 23**1 pts**

A given reaction has an activation energy of 24.52 kJ/mol. At 25°C, the half-life is 4 minutes. At what temperature will the half-life be reduced to 20 seconds?

- 150°C
- 125°C
- 115°C
- 100°C

Question 24**1 pts**

For the reaction below:



a plot of $\ln K$ vs $1/T$ gives a straight line with a slope equal to $-5.1 \times 10^3 \text{ K}$. What is the activation energy for this reaction?

- 12 kJ/mol
- 5.1 kJ/mol
- 98 kJ/mol
- 42 kJ/mol

Question 25**1 pts**

A certain reaction has an activation energy of 0.8314 kJ/mol and a rate constant of 2.718 s^{-1} at -73°C . At -173°C , which expression for the rate constant is correct?

- $\ln(k_2) = 0.5$
- $\ln(k_2) = -0.5$
- $\ln(k_2) = 1.5$
- $\ln(k_2) = 1$

Question 26**1 pts**

A food substance kept at 0°C becomes rotten (as determined by a good quantitative test) in 8.3 days. The same food rots in 10.6 hours at 30°C. Assuming the kinetics of the microorganisms enzymatic action is responsible for the rate of decay, what is the activation energy for the decomposition process? Hint: Rate varies INVERSELY with time; a faster rate produces a shorter decomposition time.

- 0.45 kJ/mol
- 23.4 kJ/mol
- 67.2 kJ/mol
- 2.34 kJ/mol

Question 27

1 pts

A catalyst...

- speeds up the reaction but does not change K.
- changes the reaction mechanism to ensure that K is increased.
- speeds up the reaction and increases K to favor product formation.
- increases K to favor product formation.

Question 28

1 pts


All else being equal, a reaction with a higher activation energy compared to one with a lower activation energy will...

- be more exothermic.
- proceed faster.
- be more endothermic.
- proceed slower.

Question 29

1 pts

Consider the potential energy diagram below:



What is the change in enthalpy (ΔH) for the reaction $A \rightarrow B$?

- 100 kJ
- 350 kJ
- 100 kJ
- 350 kJ

Question 30

1 pts

Consider the potential energy diagram in question 39. What is the activation energy (E_a) for the reaction?

- 250 kJ
- 100 kJ
- 350 kJ
- 200 kJ

Question 31

1 pts

Which of the following statements is TRUE?

- If the exponents in the rate-law do not match the coefficients in the balanced chemical equation, then we know that the reaction takes place in one step.
- The exponents in the rate-law must match the coefficients in the balanced chemical equation for the reaction.
- The rate-law for a reaction can be predicted from the balanced chemical equation.
- If the exponents in the rate-law do not match the coefficients in the balanced equation, then we know that the reaction does not take place in one step.

Question 32

1 pts

"Reaction mechanisms usually involve only unimolecular or bimolecular steps."

Is this statement true or false?

- True, because steps of higher molecularity would not be compatible with observed reaction rate laws.
- True, because the activation energy for collisions of higher molecularity would be too great.
- True, because collisions of higher molecularity would occur too infrequently to account for an observed rate.
- False.

Question 33

1 pts

Which of the following is/are ALWAYS true concerning collision and transition state theory?

I) Transition states are short-lived.

II) A balanced reaction shows which species must collide for the reaction to occur.

III) Intermediates are short-lived.

II and III

I and III

All are true.

II only





I only

III only

Question 34

1 pts

Consider the following reaction mechanism:



Which species is/are intermediates?

ClCO

Cl, ClCO

Pt, Cl, ClCO

Pt, Cl

Pt

Not saved

Submit Quiz