


This print-out should have 25 questions. Multiple-choice questions may continue on the next column or page – find all choices before answering.

001 10.0 points

How many moles of $\text{Cl}_2(\text{g})$ are produced by the electrolysis of concentrated sodium chloride if 2.00 A are passed through the solution for 4.00 hours? The equation for this process (the “chloralkali” process) is



1. 0.149 mol
2. 0.447 mol
3. 0.298 mol
4. 0.0745 mol
5. 0.00248 mol

002 10.0 points

A steel surface has been electroplated with 5.10 g of vanadium (V, molar mass = 51 g/mol). If 2.90×10^4 C of charge were used, what was the original oxidation number of V?

1. +4
2. +3
3. +5
4. +6
5. +1
6. +2

003 10.0 points

How long will it take to deposit 0.00235 moles of gold by the electrolysis of $\text{KAuCl}_4(\text{aq})$ using a current of 0.214 amperes?

1. 26.5 min

2. 17.7 min

3. 106 min

4. 70.7 min

5. 53.0 min

004 10.0 points

Consider 3 electrolysis experiments:

The first: 1 Faraday of electricity is passed through a solution of AgNO_3 .

The second: 2 Faradays of electricity are passed through a solution of $\text{Zn}(\text{NO}_3)_2$.

The third: 3 Faradays of electricity are passed through a solution of $\text{Bi}(\text{NO}_3)_3$.

1. Equal numbers of moles of all three metals are produced.

2. Twice as many moles of metallic zinc are produced than metallic silver.

3. The reaction producing the smallest mass of metal is that of the silver solution.

4. Equal masses of all three metals are produced.


005 10.0 points

Sodium is produced by electrolysis of molten sodium chloride. What are the products at the anode and cathode, respectively?

1. $\text{Cl}_2(\text{g})$ and $\text{Na}_2\text{O}(\ell)$
2. $\text{Cl}_2(\text{g})$ and $\text{Na}(\ell)$
3. $\text{Na}(\ell)$ and $\text{O}_2(\text{g})$
4. $\text{Cl}^-(\text{aq})$ and $\text{Na}_2\text{O}(\ell)$
5. $\text{O}_2(\text{g})$ and $\text{Na}(\ell)$

006 10.0 points

What is the standard cell potential of the strongest battery that could be made using these half reactions?



1. 1.11

2. 1.84

3. 1.83


4. 1.03

5. 1.04

6. 2.56

007 10.0 points

What would be the E° cell of an electrolytic cell made from the half reactions



1. -1.44


2. 1.44

3. -1.88





4. 1.88

008 (part 1 of 3) 10.0 points

The galvanic cell below uses the standard half-cells $\text{Mg}^{2+} | \text{Mg}$ and $\text{Zn}^{2+} | \text{Zn}$, and a salt bridge containing KCl(aq) . The voltmeter gives a positive voltage reading.



Identify A and write the half-reaction that occurs in that compartment.

**009 (part 2 of 3) 10.0 points**

What happens to the size of the electrode A during the operation of the cell?

1. increases

2. No change

3. decreases

010 (part 3 of 3) 10.0 points

What is the voltmeter reading?

1. +4.30 V

2. +3.40 V

3. +2.50 V

4. +0.50 V

5. +1.60 V

011 10.0 points


The electrolysis of an aqueous sodium chloride solution using inert electrodes produces

gaseous chlorine at one electrode. At the other electrode gaseous hydrogen is produced and the solution becomes basic around the electrode. What is the equation for the cathode half-reaction in the electrolytic cell?

1. $2 \text{Cl}^- \rightarrow \text{Cl}_2 + 2 e^-$
2. $2 \text{H}_2\text{O} + 2 e^- \rightarrow \text{H}_2 + 2 \text{OH}^-$
3. $\text{Cl}_2 + 2 e^- \rightarrow 2 \text{Cl}^-$
4. $\text{H}_2 + 2 \text{OH}^- \rightarrow 2 \text{H}_2\text{O} + 2 e^-$
5. None of the other answers listed is correct.

012 10.0 points


What is the E°_{cell} of



1. -0.85
2. $+1.61$
3. -2.37
4. $+2.37$
5. $+0.85$

013 10.0 points

Calculate the cell potential for a cell based on the reaction



when the concentrations are as follows: $[\text{Ag}^+] = 0.51 \text{ M}$, $[\text{Cu}^{2+}] = 0.9 \text{ M}$. (The temperature is 25°C and $E^\circ = 0.4624 \text{ V}$.)

Answer in units of V


014 10.0 points

Standard reduction potentials are established by comparison to the potential of which half reaction?

1. $\text{F}_2 + 2 e^- \rightarrow 2 \text{F}^-$
2. $\text{Li}^+ + e^- \rightarrow \text{Li}$
3. $2 \text{H}^+ + 2 e^- \rightarrow \text{H}_2$
4. $2 \text{H}_2\text{O} + 2 e^- \rightarrow \text{H}_2 + 2 \text{OH}^-$
5. $\text{Na}^+ + e^- \rightarrow \text{Na}$

015 10.0 points

Consider the cell




If E° for the cell is 0.36 V at 25°C , write the Nernst equation for the cell at this temperature.

1. $E = 0.36 - 0.01285 \ln \left[\frac{1.90}{(0.70)^2(0.60)} \right]$
2. $E = 0.36 - 0.02569 \ln \left[\frac{192.5}{(0.70)^2(0.60)} \right]$
3. $E = 0.36 - 0.01285 \ln \left[\frac{1.90}{(0.70)(0.60)} \right]$
4. $E = 0.36 + 0.01285 \ln \left[\frac{1.9}{(0.70)^2(0.60)} \right]$
5. $E = 0.36 + 0.01285 \ln \left[\frac{192.5}{(0.70)^2(0.60)} \right]$

016 10.0 points

A concentration cell consists of the same redox couples at the anode and the cathode, with different concentrations of the ions in the respective compartments. Find the unknown concentration for the following cell.



Answer in units of M

017 10.0 points

What is ratio of $[\text{Co}^{2+}] \mid [\text{Ni}^{2+}]$ when a battery built from the two half reactions



reaches equilibrium?

1. 3.20

2. 0.10

3. 0.31

4. 10.23

018 10.0 points

If E° for the disproportionation of $\text{Cu}^+(\text{aq})$ to $\text{Cu}^{2+}(\text{aq})$ and $\text{Cu}(\text{s})$ is $+0.37 \text{ V}$ at 25°C , calculate the equilibrium constant for the reaction.

1. 1.3×10^3 2. 3.2×10^{12} 3. 2.4×10^2 4. 5.7×10^{18} 5. 1.8×10^6 **019 10.0 points**

You turn on a flashlight containing brand new NiCad batteries and keep it lit for a minute or two. Which of the following can be considered TRUE regarding the chemical state of these batteries?

- I. ΔG for the battery reaction is negative.
- II. $E_{\text{cell}} > 0$.
- III. The batteries are at equilibrium.
- IV. E_{cell} is substantially decreasing during this time.

1. All

2. II and IV only

3. III only

4. All but IV


5. All but III

6. Maybe IV and I

7. I and II only

020 10.0 points

What is ΔG° for the half reaction below?

1. $194,000 \text{ kJ} \cdot \text{mol}^{-1}$ 2. $-1,418 \text{ kJ} \cdot \text{mol}^{-1}$ 3. $-709 \text{ kJ} \cdot \text{mol}^{-1}$ 4. $194 \text{ kJ} \cdot \text{mol}^{-1}$ **021 10.0 points**


For the reduction of Cu^{2+} by Zn , $\Delta G^\circ = -212 \text{ kJ/mol}$ and $E^\circ = +1.10 \text{ V}$. If the coefficients in the chemical equation for this reaction are multiplied by 2, $\Delta G^\circ = -424 \text{ kJ/mol}$. This means $E^\circ = +2.20 \text{ V}$.

1. False

2. True

022 10.0 points


Consider the cell



at standard conditions. Calculate the value of ΔG_r° for the reaction that occurs when current is drawn from this cell.

1. $-62 \text{ kJ} \cdot \text{mol}^{-1}$ 2. $+230 \text{ kJ} \cdot \text{mol}^{-1}$ 3. $-31 \text{ kJ} \cdot \text{mol}^{-1}$ 4. $+62 \text{ kJ} \cdot \text{mol}^{-1}$ 5. $-230 \text{ kJ} \cdot \text{mol}^{-1}$ **023 10.0 points**

The standard potential of the cell



is +0.23 V at 25°C. Calculate the equilibrium constant for the reaction of 1 M $\text{Pb}^{2+}(\text{aq})$ with 1 M $\text{SO}_4^{2-}(\text{aq})$.

1. 3.7×10^{16}

2. 1.7×10^{-8}


3. 8.0×10^{17}

4. 6.0×10^7

5. 7.7×10^3

024 10.0 points

The standard voltage of the cell



is +0.73 V at 25°C. Calculate the equilibrium constant for the cell reaction.

1. 2.2×10^{12}

2. 4.6×10^{-13}


3. 5.1×10^{14}

4. 2.0×10^{-15}

5. 3.9×10^{-29}

025 10.0 points

The equilibrium constant for the reaction



is 5.6×10^{-20} at 25°C. Calculate the value of E° for a cell utilizing this reaction.

1. + 1.14 V

2. – 0.57 V

3. + 0.57 V

4. – 1.14 V

5. – 0.25 V