

last name

first name

signature



REMEMBER: Bubble in ALL Bubblesheet information!

This includes your first and last name, your UTEID, and your version number.

Please refer to the back of the bubble sheet for more info.

$R = 8.314 \text{ J/mol}\cdot\text{K}$	PV = nRT
R=0.08206L atm/mol·K	
$R = 62.36 \text{ L torr/mol} \cdot \text{K}$	$q = m \cdot C_{\rm s} \cdot \Delta I$ $q = m \cdot \Delta H_{\rm change}$
$1 \text{ atm} = 1.01325 \times 10^5 \text{ Pa}$	$\ln\left(\frac{P_2}{P_1} ight) = rac{\Delta H_{ ext{vap}}}{R}\left(rac{1}{T_1} - rac{1}{T_2} ight)$
1  atm = 760  torr	$\Delta H_{\rm solution} = \Delta H_{\rm lattice} + \Delta H_{\rm hydration}$
1  atm = 14.7  psi	
	$P_{\rm A} = \chi_{\rm A} \cdot P_{\rm A}^{\rm o}$
water data	
$K_{\mathrm{f}} = 1.86~^{\circ}\mathrm{C}/m$	$\Delta T_{\rm f} = i \cdot K_{\rm f} \cdot m \qquad \Delta T_{\rm b} = i \cdot K_{\rm b} \cdot m$
$K_{\rm b}=0.512~{\rm ^{\circ}C}/m$	$\Pi = i \cdot MRT$
$C_{\rm s,ice} = 2.09~{\rm J/g~K}$	
$C_{\rm s,water} = 4.184 \text{ J/g K}$	$G = H - TS$ $\Delta G = \Delta H - T\Delta S$
$C_{\rm s,steam} = 2.03~\rm J/g~K$	
$\Delta H_{\rm fus} = 334 \ {\rm J/g}$	
$\Delta H_{\mathrm{vap}} = 2260 \mathrm{~J/g}$	

**NOTE:** Please keep your Exam copy intact (all pages still stapled). You must turn in your exam copy, plus your bubble sheet, and any scratch paper.

This print-out should have 25 questions. Multiple-choice questions may continue on the next column or page – find all choices before answering.

#### 001 4.0 points

On a hiking expedition with Bear Grylls, you 'accidentally' end up in a huge chasm 2 km below sea level. Bear, being the resourceful TV star that he is, builds a fire and promptly starts to heat some water to cook the local snake for dinner. At what temperature do you expect Bear's water to boil?

**1.** At  $100^{\circ}$ C exactly

**2.** Higher than 100°C **correct** 

**3.** None of these; Bear Grylls eats his food raw!

**4.** Lower than  $100^{\circ}$ C

#### **Explanation**:

At sea level, the pressure is 1 atm (760 torr). As you decrease elevation, pressure increases, raising the boiling point of water.

#### **002 4.0** points

Nitrogen gas,  $N_2(g)$ , has a certain solubility when dissolved in water. In which of the following cases would the solubility of  $N_2(g)$ increase?

I) changing to a less polar solvent

- II) increasing the amount of solvent
- III) increasing the pressure of  $N_2(g)$

1. I only

2. I, II and III

**3.** I and II

4. I and III correct

5. II only

6. III only

7. II and III

### Explanation:

The solubility of a gas is directly proportional to the gas' pressure and inversely proportional to the temperature of the solvent. Considering the principle of like dissolve like, a non-polar species such as  $N_2(g)$  will be more soluble in non-polar solvents. The volume of the solvent does not impact the solubility, which is an intensive property; this is evident when one considers the units of solubility such as grams per liter, etc.

### 003 4.0 points

A solution containing all of the solute that a solvent can dissolve at a certain temperature and pressure is called

**1.** an unsaturated solution.

**2.** a supersaturated solution.

**3.** a concentrated solution.

4. a saturated solution. correct

### Explanation:

### **004 4.0** points

What is  $K_{\rm sp}$  for Ag<sub>3</sub>PO<sub>4</sub>, if its molar solubility is  $2.7 \times 10^{-6}$  mol/L?

r of N<sub>2</sub>(g) 1.  $1.4 \times 10^{-21}$  correct 2.  $7.3 \times 10^{-12}$ 3.  $5.3 \times 10^{-16}$ 4.  $2.0 \times 10^{-17}$ 5.  $5.3 \times 10^{-23}$ 6.  $4.8 \times 10^{-22}$ 7.  $1.7 \times 10^{-14}$ Explanation:  $S = 2.7 \times 10^{-6}$  mol/L The solubility equilibrium is

$$Ag_3PO_4(s) \rightleftharpoons 3Ag^+(aq) + PO_4^{3-}(aq)$$

$$[Ag^+] = 3 S = 8.1 \times 10^{-6} \text{ mol/L} [PO_4^{3-}] = S = 2.7 \times 10^{-6} \text{ mol/L}$$

$$K_{\rm sp} = [{\rm Ag}^+]^3 [{\rm PO}_4^{3-}]$$
  
=  $(8.1 \times 10^{-6})^3 (2.7 \times 10^{-6})$   
=  $1.43489 \times 10^{-21}$ 

# 005 4.0 points

The phase diagram for a pure compound is given below.



All of the following could have a similar phase diagram except

**1.** methanol.

2. carbon dioxide.

3. benzene.

4. water. correct

5. carbon tetrachloride.

## **Explanation:**

# 006 4.0 points

Theoretically, it would be harder to dissolve  $(NaCl/Al_2S_3)$  in water because the (higher/lower) the charge density, the lower the solubility.

- 1.  $Al_2S_3$ , higher correct
- **2.**  $Al_2S_3$ , lower

3. NaCl, higher

4. NaCl, lower

# Explanation:

 $Al_2S_3$  has a higher charge density  $(Al^{3+}, S^{2-})$  than NaCl (Na<sup>+</sup>, Cl<sup>-</sup>). High charge density corresponds to high lattice energy and thus much more endothermic (i.e. less spontaneous) dissolution.

# 007 4.0 points

What mass of ethylene glycol  $((CH_2OH)_2)$ with molecular weight 62 g/mol) must be added to 1.00 L of H<sub>2</sub>O (of mass 1 kg) to lower the freezing point to  $-5^{\circ}C$ ?  $K_{\rm f H_2O} =$  $1.86^{\circ}C/m$ .

**1.** 330 g

**2.** 66 g

**3.** 25 g

4. 167 g correct

**5.** 123 g

# Explanation:

 $\begin{aligned} \mathrm{MW}_{\mathrm{CH}_{2}\mathrm{OH}} &= 62 \text{ g/mol} & V_{\mathrm{H}_{2}\mathrm{O}} &= 1 \text{ L} \\ \mathrm{m}_{\mathrm{H}_{2}\mathrm{O}} &= 1 \text{ kg} & T_{\mathrm{f}} &= -5^{\circ}\mathrm{C} \end{aligned}$ 

$$T_{\rm f} = T_{\rm f}^0 - \Delta T_{\rm f}$$
$$\Delta T_{\rm f} = T_{\rm f}^0 - T_{\rm f}$$
$$= 0.00^{\circ} \rm C - (-5^{\circ} \rm C)$$
$$= 5^{\circ} \rm C$$

$$\Delta T_{\rm f} = K_{\rm f} \cdot m$$
$$m = \frac{\Delta T_{\rm f}}{K_{\rm f}}$$
$$= \frac{5^{\circ} \rm C}{1.86^{\circ} \rm C/m} = 2.688 \ m$$

$$m = \frac{n_{(CH_2OH)_2}}{\text{kg water}} = \frac{\frac{g_{(CH_2OH)_2}}{MW_{(CH_2OH)_2}}}{\text{kg water}}$$

$$g (CH_2OH)_2$$

$$= m(MW_{(CH_2OH)_2})(kg water)$$

$$= \frac{2.688 \text{ mol } (CH_2OH)_2}{kg \text{ water}}$$

$$\times \frac{62.0 \text{ g } (CH_2OH)_2}{\text{mol } (CH_2OH)_2}$$

$$\times (1 \text{ kg water})$$

$$= 167 \text{ g } (CH_2OH)_2$$

### 008 4.0 points

The phase diagram for a pure substance is given below.



The substance is stored in a container at 150 atm at  $25^{\circ}$ C. Describe what happens if the container is opened at  $25^{\circ}$ C.

**1.** The liquid in the container vaporizes. **correct** 

- 2. The solid in the container melts.
- **3.** The vapor in the container escapes.
- 4. The liquid in the container freezes.
- 5. The solid in the container sublimes.

### **Explanation:**

**009 4.0 points**  $\Delta G_{\text{vap}}^{\circ}$  for H<sub>2</sub>O( $\ell$ ) at 85°C is (< 0, = 0, > 0) and at 100°C is (< 0, = 0, > 0).

1. < 0; < 0

2. < 0; > 0

### 010 4.0 points

Estimate the enthalpy of vaporization of  $CCl_4$  given that at 25°C and 58°C its vapor pressure is 107 and 405 torr, respectively. Assume that the enthalpy of vaporization is independent of the temperature.

- **1.**  $48.6 \text{ kJ} \cdot \text{mol}^{-1}$
- **2.** 33.1 kJ·mol<sup>-1</sup> correct
- **3.**  $142 \text{ kJ} \cdot \text{mol}^{-1}$
- **4.**  $3.98 \text{ kJ} \cdot \text{mol}^{-1}$
- **5.**  $486 \text{ J} \cdot \text{mol}^{-1}$

## Explanation: $T_1 = 25^{\circ}\text{C} + 273.15 = 298.15 \text{ K}$

 $P_1 = 107 \text{ torr}$  $T_2 = 58^{\circ}\text{C} + 273.15 = 331.15 \text{ K}$  $P_2 = 405 \text{ torr}$ Using the Clausius-Clapeyron equation,

$$\ln\left(\frac{P_{2}}{P_{1}}\right) = \frac{\Delta H_{\text{vap}}^{\circ}}{R} \left(\frac{1}{T_{1}} - \frac{1}{T_{2}}\right)$$
$$\Delta H_{\text{vap}}^{0} = \frac{R \ln\left(\frac{P_{2}}{P_{1}}\right)}{\frac{1}{T_{1}} + \frac{1}{T_{2}}}$$
$$= \frac{8.314 \text{ J} \cdot \text{mol}^{-1} \cdot \text{K}}{\frac{1}{298.15 \text{ K}} - \frac{1}{331.15 \text{ K}}} \times \ln\left(\frac{405 \text{ torr}}{107 \text{ torr}}\right)$$
$$= 33109 5 \text{ J} \cdot \text{mol}^{-1}$$

What is the molar solubility of Ag<sub>2</sub>S? The  $K_{\rm sp}$  is  $6.3 \times 10^{-51}$ .

1.  $6.37 \times 10^{-15}$ 

**2.**  $7.94 \times 10^{-26}$ 

**3.**  $5.8 \times 10^{-18}$ 

**4.**  $2.82 \times 10^{-13}$ 

**5.**  $1.16 \times 10^{-17}$  correct

### **Explanation:**

#### 012 4.0 points

Consider a 200 g block of ice at standard pressure. If it is initially at -23 °C and is heated until it is steam at 148 °C, how much total heat was added to the sample of water? Use the following thermodynamic values for your calculation:

$$c_{ice} = 2.09 \text{ J} \cdot \text{g}^{-1} \cdot \text{K}^{-1}$$

$$c_{water} = 4.184 \text{ J} \cdot \text{g}^{-1} \cdot \text{K}^{-1}$$

$$c_{steam} = 2.03 \text{ J} \cdot \text{g}^{-1} \cdot \text{K}^{-1}$$

$$\Delta H_{vap} = 2260 \text{ J} \cdot \text{g}^{-1}$$

$$\Delta H_{fus} = 334 \text{ J} \cdot \text{g}^{-1}$$

1.565 kJ

**2.** 548 kJ

**3.** 822 kJ

4.29.1 kJ

**5.** 632 kJ correct

#### **Explanation**:

 $\Delta T_1 = 0^{\circ} \text{C} - -23 \,^{\circ} \text{C} = 23 \,^{\circ} \text{C}$   $\Delta T_2 = 100^{\circ} \text{C}$   $\Delta T_3 = 148 \,^{\circ} \text{C} - 100^{\circ} \text{C} = 48 \,^{\circ} \text{C}$  m = 200 gq = 632 kJ

## 013 4.0 points

Suppose that you wanted to be sure that a metal ion, any metal ion, would dissolve in water.

What salt of the metal ion compound would you choose?

**1.** the nitrate  $(NO_3^-)$  salt of the metal ion **correct** 

**2.** the hydroxide  $(OH^-)$  salt of the metal ion

**3.** the carbonate  $(CO_3^{2-})$  salt of the metal ion

**4.** the chloride  $(Cl^{-})$  salt of the metal ion

#### Explanation:

Nitrate  $(NO_3^-)$  salts of any metal are always soluble.

### 014 4.0 points

Which of the following salts would have the greatest molar solubility?

- **1.** Al(OH)<sub>3</sub>  $K_{sp} = 1.90 \times 10^{-33}$
- **2.** PbCrO<sub>4</sub>  $K_{sp} = 1.77 \times 10^{-14}$  correct
- **3.** CdS  $K_{sp} = 3.60 \times 10^{-29}$
- 4. Cu<sub>2</sub>S  $K_{sp} = 2.00 \times 10^{-47}$

#### Explanation:

Lead chromate is a salt composed of only two ions and thus its molar solubility is equal to the square root of its  $K_{sp}$ . The best approximation is therefore  $(10^{-14})^{1/2} = 10^{-7}$ .

#### 015 4.0 points

Which of the following highly soluble salts would be the most useful for lowering the freezing point of a solution?

**1.**  $Ce_2(SeO_4)_3$  correct

**2.**  $(NH_4)_3PO_4$ 

**3.** KBr

4.  $Cs_2SO_4$ 

5.  $Cs_2Cr_2O_7$ 

#### **Explanation:**

Cerium Selenate,  $Ce_2(SeO_4)_3$ , has the largest van't Hoff factor (i = 5), and would require the smallest molar quantity to lower a solution's freezing point by a given amount.

#### 016 4.0 points

Water from a local stream is added to one side of the U-tube shown below. Pure water is placed in the tube on the other side of the semipermeable membrane. With the left side open to barometric pressure of 1.0 atm and 1.15 atm applied to the right side, the two liquids do not move.



In which half of the U-tube is the pure water located?

#### 1. A correct

**2.** B

**3.** Not enough information is given.

### **Explanation:**

### 017 4.0 points

In general, decreasing the temperature makes which phase transitions more likely to occur?

1. evaporation, fusion, sublimation

2. evaporation, deposition, freezing

**3.** sublimation, condensation, freezing

4. condensation, freezing, deposition correct

5. condensation, fusion, deposition

# Explanation:

Phase changes are equilibrium processes and are thus temperature dependent. Also, the  $\Delta$ H and  $\Delta$ S always have the same sign. Lowering the temperature makes exothermic phase transitions more likely - freezing, condensation and deposition.

# 018 4.0 points

Pure water is saturated with  $PbCl_2$ . In this saturated solution

[Pb<sup>2+</sup>] = 0.5 [Cl<sup>-</sup>]. correct
 K<sub>sp</sub> = [Pb<sup>2+</sup>].
 [Pb<sup>2+</sup>] [Cl<sup>-</sup>] = K<sub>sp</sub>.
 [Pb<sup>2+</sup>] = [Cl<sup>-</sup>].
 2 [Cl<sup>-</sup>] = [Pb<sup>2+</sup>].
 Explanation:

### 019 4.0 points

Assume the molar solubility of silver chromate  $(Ag_2CrO_4)$  is represented as x. Which of the following expressions correctly expresses the relationship between the molar solubility of silver chromate and the solubility product constant  $(K_{sp})$  for this compound?

**1.** 
$$K_{sp} = 2 x^2$$
  
**2.**  $K_{sp} = x^2$ 

**3.** 
$$K_{\rm sp} = 4 x^2$$

**4.**  $K_{\rm sp} = 2 x^3$ 

5.  $K_{\rm sp} = 4 x^3$  correct

# **Explanation:**

The equation for the dissociation of  $Ag_2CrO_4$  is written as

$$Ag_2CrO_4 \rightleftharpoons 2Ag^+ + CrO_4^-$$

The expression for  $K_{\rm sp} = \left[ {\rm Ag}^+ \right]^2 \left[ {\rm CrO}_4^- \right]$ . Since the molar solubility is x,  $K_{\rm sp}$  can be rewritten (in terms of x) as

$$K_{\rm sp} = [2 x]^2 [x] = 4 x^3$$
.

 $[Ag^+] = 2x$  because for every salt molecule that dissociates,  $2Ag^+$  ions are produced.

# 020 4.0 points

The vapor pressure of pure  $CH_2Cl_2$  (with molecular weight 85 g/mol) is 133 torr at 0°C and the vapor pressure of pure  $CH_2Br_2$ (with molecular weight 174 g/mol) is 11 torr at the same temperature. What is the total vapor pressure at 0°C of a solution prepared from equal masses of these two substances?

**1.** 124 torr

**2.** 89 torr

- **3.** 72 torr
- **4.** 7.4 torr
- 5. 3.6 torr
- **6.** 105 torr
- 7.93 torr correct
- **8.** 144 torr
- **9.** 44 torr

### **Explanation:**

For CH<sub>2</sub>Cl<sub>2</sub>,  $P^0 = 133 \text{ torr}$  MW = 85 g/mol For CH<sub>2</sub>Br<sub>2</sub>,  $P^0 = 11 \text{ torr}$  MW = 174 g/mol The mole fractions are

$$X_{\rm CH_2Cl_2} = \frac{n_{\rm CH_2Cl_2}}{n_{\rm CH_2Cl_2} + n_{\rm CH_2Cl_2}}$$
$$= \frac{g/85}{g/85 + g/174} = 0.672$$

$$X_{\rm CH_2Br_2} = \frac{n_{\rm CH_2Br_2}}{n_{\rm CH_2Br_2} + n_{\rm CH_2Cl_2}}$$
$$= \frac{g/174}{g/174 + g/85} = 0.328$$

Thus

$$P_{\rm solution} = P_{\rm CH_2Cl_2} + P_{\rm CH_2Br_2}$$

 $= X_{\text{CH}_2\text{Cl}_2} P^0_{\text{CH}_2\text{Cl}_2}$  $+ X_{\text{CH}_2\text{Br}_2} P^0_{\text{CH}_2\text{Br}_2}$ = (0.672)(133 torr)+ (0.328)(11 torr)= 93.0 torr

# **021 4.0** points

Rank the liquids

NH<sub>3</sub>, CH<sub>3</sub>OH, CH<sub>3</sub>CH<sub>2</sub>F, CCl<sub>4</sub> by their miscibility in heptane ( $C_7H_{16}$ ), from most miscible to least.

NH<sub>3</sub> > CH<sub>3</sub>OH > CH<sub>3</sub>CH<sub>2</sub>F > CCl<sub>4</sub>
 CCl<sub>4</sub> > CH<sub>3</sub>CH<sub>2</sub>F > NH<sub>3</sub> > CH<sub>3</sub>OH
 CH<sub>3</sub>CH<sub>2</sub>F > CCl<sub>4</sub> > CH<sub>3</sub>OH > NH<sub>3</sub>
 CH<sub>3</sub>CH<sub>2</sub>F > CH<sub>3</sub>OH > CCl<sub>4</sub> > NH<sub>3</sub>
 CCl<sub>4</sub> > CH<sub>3</sub>CH<sub>2</sub>F > CH<sub>3</sub>OH > NH<sub>3</sub>
 CCl<sub>4</sub> > CH<sub>3</sub>CH<sub>2</sub>F > CH<sub>3</sub>OH > NH<sub>3</sub>

#### **Explanation:**

"Like dissolves like." Heptane is hydrocarbon, and thus completely non-polar. Carbon tetrachloride is most soluble since even though the C-Cl bond is polar, there is no net dipole in the molecule, followed by fluoroethane, which is polar. Methanol is small and very polar (exhibiting hydrogen bonding), and ammonia is the most polar of the liquids, and thus least soluble.

### 022 4.0 points

 $K_{\rm sp}$  for CaF<sub>2</sub> is  $3.9 \times 10^{-11}$ . Would a precipitate of CaF<sub>2</sub> form if Ca(NO<sub>3</sub>)<sub>2</sub> and NaF solutions were mixed such that [Ca<sup>2+</sup>] =  $2.0 \times 10^{-4}$  M, and [F<sup>-</sup>] =  $3.0 \times 10^{-4}$  M?

**1.** yes, because Q is larger than  $K_{\rm sp}$ 

**2.** yes, because Q is smaller than  $K_{\rm sp}$ 

3. no correct

# Explanation:

023 4.0 points

Identify the spectator ion(s) in the equation  $CaCl_2(aq) + Na_2CO_3(aq) \rightarrow$  $CaCO_3(s) + 2NaCl(aq).$ 

- 1.  $Na^+$ ,  $Cl^-$  correct
- **2.**  $Ca^{2+}$ ,  $CO_3^{2-}$
- **3.**  $Ca^{2+}$ ,  $Cl^{-}$

4. Na<sup>+</sup>,  $CO_3^{2-}$ 

# Explanation:

Spectator ions are those that do not participate in the reaction nor appear in the net ionic equation. Sodium ions  $(Na^+)$ , potassium ions  $(K^+)$ , nitrate  $(NO_3^-)$  and chloride  $(Cl^-)$  are common spectator ions.

### 024 4.0 points

The solubility product constant of Ag<sub>2</sub>CrO<sub>4</sub> is  $9.0 \times 10^{-12}$ . What is the molar solubility of Ag<sub>2</sub>CrO<sub>4</sub> in a solution in which the silver ion concentration is maintained at  $2.0 \times 10^{-3}$  M by addition of AgNO<sub>3</sub>?

**1.**  $5.6 \times 10^{-7}$ 

**2.**  $4.0 \times 10^{-3}$ 

**3.**  $2.3 \times 10^{-6}$  correct

**4.**  $1.3 \times 10^{-4}$ 

**5.**  $4.5 \times 10^{-9}$ 

# **Explanation:**

### 025 4.0 points

Which of the following would raise the vapor pressure of a sample of isopropanol in a closed container?

- I) increasing the size of the container
- II) increasing the temperature of the sample
- III) increasing the applied pressure

## 1. III only

2. I, II and III

I only
 I and III
 I and II
 I and II
 II and III

# 7. II only correct

## Explanation:

Because evaporation is always an endothermic process, a substances vapor pressure is always directly proportional to its temperature. Adding a non-volatile solute to any substance will decrease its vapor pressure. Factors such as sample size and container size do not influence he vapor pressure.