

# HW07 - Kinetics

Started: Mar 22 at 10:56am

## Quiz Instructions

### Homework 07

### Kinetics

#### Question 1

0.5 pts

Consider the reaction:



What is the overall order of the reaction and the order with respect to  $[\text{O}_3]$ ?

1 and 2

2 and 2


-1 and 3

3 and 2

#### Question 2

0.5 pts

When the reaction below:



is proceeding under conditions such that 0.015 mol/L of  $\text{N}_2\text{O}$  is being formed each second, the rate of the overall reaction is \_\_\_\_\_ and the rate of change for NO is \_\_\_\_\_.

0.015 M/s; -0.005 M/s

0.015 M/s; -0.045 M/s


0.030 M/s; -0.005 M/s

0.015 M/s; 0.045 M/s

### Question 3

1 pts

What is the rate law for the reaction below:



if the following data were collected?


Exp	[A] <sub>0</sub>	[B] <sub>0</sub>	[C] <sub>0</sub>	Initial Rate
1	0.4	1.2	0.7	2.32x10 <sup>-3</sup>
2	1.3	1.2	0.9	7.54x10 <sup>-3</sup>
3	0.4	4.1	0.8	9.25x10 <sup>-2</sup>
4	1.3	1.2	0.2	7.54x10 <sup>-3</sup>

- rate = 1.49x10<sup>-3</sup> [B]<sup>3</sup> [C]
- rate = 5.37x10<sup>-3</sup> [A] [B]<sup>3</sup>
- rate = 1.79x10<sup>-3</sup> [B]<sup>2</sup> [C]
- rate = 3.36x10<sup>-3</sup> [A] [B]<sup>3</sup>
- rate = 4.48x10<sup>-3</sup> [A] [B]<sup>2</sup> [C]

### Question 4

1 pts

A chemical reaction is expressed by the balanced chemical equation:



Consider the data below:

exp	[A] <sub>0</sub>	[B] <sub>0</sub>	initial rate (M/min)
1	0.15	0.15	0.00110363
2	0.15	0.3	0.0044145
3	0.3	0.3	0.008829

Find the rate law for the reaction.

rate = k [A] [B]

rate = k [A] [B]<sup>2</sup>

rate = k [A]<sup>2</sup> [B]

rate = k [B]<sup>2</sup>

**Question 5**

0.5 pts

Calculate the value of the rate constant (k) for the reaction in question 4.

0.327

0.000166

0.00110

0.00736

**Question 6**

0.5 pts

If the initial concentrations of both A and B are 0.31 M for the reaction in questions 4 and 5, at what initial rate is C formed?

0.00974 M/min

0.0314 M/min


0.101 M/min

-0.00974 M/min

**Question 7**

1 pts

We know that the rate expression for the reaction below:



at a certain temperature is  $\text{rate} = [\text{NO}]^2 [\text{O}_2]$ . We carry out two experiments involving this reaction at the same temperature, but in the second experiment the initial concentration of NO is doubled while the initial concentration of  $\text{O}_2$  is halved. The initial rate in the second experiment will be how many times that of the first?

8

1

2

4

### Question 8

1 pts

Consider the data collected for a chemical reaction between compounds A and B that is first order in A and first order in B:

rxn	[A] <sub>0</sub>	[B] <sub>0</sub>	rate (M/s)
1	0.2	0.05	0.1
2	?	0.05	0.4
3	0.4	?	0.8

From the information above for 3 experiments, determine the missing concentrations of A and B. Answers should be in the order [A] then [B].

0.20 M; 0.80 M

0.40 M; 0.10 M

0.40 M; 0.20 M

0.80 M; 0.10 M

0.80 M; 0.20 M

### Question 9

0.5 pts


For a reaction that is zero-order overall...

the reactant concentration does not change with time.

- the activation energy is zero.
- the rate does not change during the reaction.
- the rate constant is zero.

**Question 10****0.5 pts**

Consider the reaction below:




If it is 1st order in A and 0th order in B, a plot of  $\ln[A]$  vs time will have a slope that is...

- decreasing exponentially.
- increasing exponentially.
- constant.
- slowly increasing.

**Question 11****1 pts**

Consider the reaction below:




If it has a half-life of 1.6 sec, how long will it take a system with  $[\text{H}_2\text{CO}_3]_0$  of 2M to reach  $[\text{H}_2\text{CO}_3]$  of 125mM?

- 2.9 sec
- 3.2 sec
- Not enough information is given.
- 6.4 sec

**Question 12****1 pts**

At a certain fixed temperature, the reaction below:




is found to be first order in the concentration of A and zeroth order in the concentration of B. The reaction rate constant is  $0.05\text{s}^{-1}$ . If 2.00 moles of A and 4.00 moles of B are placed in a 1.00 liter container, how many seconds will elapse before the concentration of A has fallen to 0.30 moles/liter?

- There is not enough information to answer.
- 37.94 sec
- 10.22 sec
- 2.83 sec

### Question 13

0.5 pts

The reaction below:




is observed to obey first-order kinetics. Which of the following plots should give a straight line?

- $\ln[A]$  vs  $k$
- $[A]$  vs  $k$
- $\ln[A]$  vs  $k^{-1}$
- $\ln[A]$  vs  $t^{-1}$
- $[A]$  vs  $t^{-1}$
- $[A]$  vs  $t$
- $\ln[A]$  vs  $t$

### Question 14

1 pts

For the reaction below:



at 800K, a plot of  $\ln[\text{cyclobutane}]$  vs  $t$  gives a straight line with a slope of  $-1.6\text{ s}^{-1}$ . Calculate the time needed for the

concentration of cyclobutane to fall to 1/16 of its initial value.

- 1.7 sec
- 0.63 sec
- 1.6 sec
- 1.3 sec

**Question 15**

1 pts

The initial concentration of the reactant A in a first-order reaction is 1.2 M. After 69.3 sec, the concentration has fallen to 0.3 M. What is the rate constant  $k$ ?

- not enough information
- $0.01 \text{ s}^{-1}$
- $0.02 \text{ s}^{-1}$
- $0.2 \text{ s}^{-1}$

**Question 16**

1 pts


A reaction is found to be first order with respect to one of the reactant species, A. When might a plot of  $\ln[A]$  vs time NOT yield a straight line?

- when the rate also depends on the concentration of another reactant as well
- All of the other answers could be correct.
- if the reaction comes to equilibrium
- if the reaction has any significant backward rate

**Question 17**

0.5 pts

Consider the following elementary reactions:




Identify the molecularity of each reaction respectively.

- tetramolecular, termolecular, pentamolecular
- bimolecular, unimolecular, termolecular
- all three elementary reactions are bimolecular
- it is impossible to know without knowing the overall reaction for each

### Question 18

1 pts

A and B react to form C according to the single step reaction below:




Which of the following is the correct rate equation for [B] and the correct units for the rate constant of this reaction?

- $\frac{\Delta [B]}{\Delta t} = -\frac{2k [A] [B]}{[C]}; \frac{1}{M \cdot s}$
- $\frac{\Delta [B]}{\Delta t} = -k [A] [B]^2; \frac{1}{M^2}$
- $\frac{\Delta [B]}{\Delta t} = -2k [A] [B]^2; \frac{1}{M^2 \cdot s}$
- $\frac{\Delta [B]}{\Delta t} = -2k [A] [B]; \frac{1}{M \cdot s}$


### Question 19

1 pts

Consider the mechanism below:







What is the rate law?

rate =  $k_2[\text{NO}_2]^2$

rate =  $k_1k_2[\text{NO}_2]^2$

rate =  $k_1[\text{NO}_2][\text{F}_2]$



rate =  $k_2[\text{NO}_2][\text{F}]$

rate =  $k_1[\text{NO}_2\text{F}][\text{F}_2]$

### Question 20

1 pts

Determine the overall balanced equation for a reaction having the following proposed mechanism:



and write an acceptable rate law.

$\text{A} + \text{B}_2 \longrightarrow \text{C}_2 + \text{D}$ ; rate =  $k[\text{B}_2]^2$

$\text{E}_3 + \text{A} \longrightarrow \text{B}_2 + \text{C}_2$ ; rate =  $k[\text{E}_3][\text{A}]$


$2\text{B}_2 \longrightarrow \text{E}_3 + \text{D}$ ; rate =  $k[\text{B}_2]^2$

$\text{A} + \text{B}_2 \longrightarrow \text{C}_2 + \text{D}$ ; rate =  $k[\text{A}][\text{B}_2]$



### Question 21

1 pts

Consider the reaction below:



The proposed mechanism of this reaction is:



What is the rate of the overall reaction?

$rate = k_2 [I]^2 [H_2]$

$rate = \frac{k_1 k_2}{k_{-1}} [I_2] [H_2]$

$rate = k_1 k_2 [I_2] [H_2]$

$rate = \frac{k_{-1} k_2}{k_1} [I_2] [H_2]$

$rate = \frac{k_1 k_2}{k_{-1}} [I]^2 [H_2]$

**Question 22**

1 pts

A reaction rate increases by a factor of 655 in the presence of a catalyst at 37°C. The activation energy of the original pathway is 106 kJ/mol. What is the activation energy of the new pathway, all other factors being equal?

16,600 J/mol

16,600 kJ/mol

89.3 kJ/mol

89.3 J/mol

**Question 23**

1 pts

A given reaction has an activation energy of 24.52 kJ/mol. At 25°C, the half-life is 4 minutes. At what temperature will the half-life be reduced to 20 seconds?

150°C

125°C

100°C

115°C

**Question 24**

1 pts

For the reaction below:



a plot of  $\ln K$  vs  $1/T$  gives a straight line with a slope equal to  $-5.1 \times 10^3$  K. What is the activation energy for this reaction?

42 kJ/mol

12 kJ/mol

5.1 kJ/mol

98 kJ/mol

**Question 25**

1 pts

A certain reaction has an activation energy of 0.8314 kJ/mol and a rate constant of  $2.718 \text{ s}^{-1}$  at  $-73^\circ\text{C}$ . At  $-173^\circ\text{C}$ , which expression for the rate constant is correct?

$\ln(k_2) = -0.5$

$\ln(k_2) = 0.5$

$\ln(k_2) = 1$

$\ln(k_2) = 1.5$

**Question 26**

1 pts

A food substance kept at  $0^\circ\text{C}$  becomes rotten (as determined by a good quantitative test) in 8.3 days. The same food rots in 10.6 hours at  $30^\circ\text{C}$ . Assuming the kinetics of the microorganisms enzymatic action is responsible for the rate of decay, what is the activation energy for the decomposition process? Hint: Rate varies INVERSELY with time; a faster rate produces a shorter decomposition time.

23.4 kJ/mol

0.45 kJ/mol

2.34 kJ/mol

67.2 kJ/mol

**Question 27**

1 pts

A catalyst...

- changes the reaction mechanism to ensure that K is increased.
- increases K to favor product formation.
- speeds up the reaction but does not change K.
- speeds up the reaction and increases K to favor product formation.

**Question 28**

1 pts


All else being equal, a reaction with a higher activation energy compared to one with a lower activation energy will...

- be more exothermic.
- proceed slower.
- proceed faster.
- be more endothermic.

**Question 29**

1 pts

Consider the potential energy diagram below:



What is the change in enthalpy ( $\Delta H$ ) for the reaction  $A \rightarrow B$ ?

- 350 kJ
- 100 kJ
- 100 kJ
- 350 kJ

**Question 30**

1 pts

Consider the potential energy diagram in question 39. What is the activation energy ( $E_a$ ) for the reaction?

- 250 kJ
- 350 kJ
- 100 kJ
- 200 kJ

**Question 31**

1 pts

Which of the following statements is TRUE?

- If the exponents in the rate-law do not match the coefficients in the balanced chemical equation, then we know that the reaction takes place in one step.
- The rate-law for a reaction can be predicted from the balanced chemical equation.
- The exponents in the rate-law must match the coefficients in the balanced chemical equation for the reaction.
- If the exponents in the rate-law do not match the coefficients in the balanced equation, then we know that the reaction does not take place in one step.

**Question 32**

1 pts

"Reaction mechanisms usually involve only unimolecular or bimolecular steps."

Is this statement true or false?

- True, because steps of higher molecularity would not be compatible with observed reaction rate laws.
- True, because the activation energy for collisions of higher molecularity would be too great.
- False.
- True, because collisions of higher molecularity would occur too infrequently to account for an observed rate.

**Question 33**

1 pts

Which of the following is/are ALWAYS true concerning collision and transition state theory?

I) Transition states are short-lived.

II) A balanced reaction shows which species must collide for the reaction to occur.

III) Intermediates are short-lived.




- All are true.
- III only
- I and III
- II only
- I only

II and III

**Question 34**

**1 pts**

Consider the following reaction mechanism:



Which species is/are intermediates?

Pt, Cl, ClCO

Pt

Pt, Cl

Cl, ClCO

ClCO

Quiz saved at 10:56am

Submit Quiz