HW04 - Electromagnetic Radiation

▲ This is a preview of the published version of the quiz

Started: Jul 7 at 9:43am

Quiz Instructions

Homework 04 - Electromagnetic Radiation

Question 1	1 pts
What is the frequency of light with a wavelength of 4.0×10^{-7} m?	
◯ 3.0 x 10 ⁻¹⁴ s ⁻¹	
◯ 3.0 x 10 ¹⁴ s ⁻¹	
◯ 1.3 x 10 ⁻¹⁵ s ⁻¹	
○ 7.5 x 10 ¹⁴ s ⁻¹	

Question 2	1 pts
What is the correct order of increasing frequency?	
O radio waves, visible light, ultraviolet radiation, infrared radiation	
O infrared radiation, radio waves, visible light, ultraviolet radiation	
O radio waves, infrared radiation, ultraviolet radiation, visible light	
O ultraviolet radiation, visible light, infrared radiation, radio waves	
O radio waves, infrared radiation, visible light, ultraviolet radiation	

Light with a frequency of 7.30 x 10 ¹⁴ Hz lies in the violet region of the visible spectrum. What is the wavelength of this frequency of light?	
O 411 nm	
◯ 4.11 x 10 ⁻⁷ nm	
◯ 4.11 x 10 ⁻¹⁶ nm	
○ 4.11 x 10 ²¹ nm	

Question 4	1 pts
When an electron beam strikes a block of copper, x-rays of frequency 1.97 x 10 ¹⁹ Hz are emitted. What is the of these x-rays?	e wavelength
◯ 1.52 x 10 ⁻² pm	
◯ 1.52 x 10 ⁻¹¹ nm	
○ 15.2 pm	
○ 15.2 nm	

Question 5	1 pts
Wavelength is	
O the distance between successive peaks in a wave.	
O the distance between a peak of one wave and the trough of the next.	
O one-half of the height of a wave.	
O the number of waves passing a fixed point in one second.	

Frequency is...

O one half the height of the wave.

O the distance between a peak in one wave to the trough in the next wave.

the number of waves passing a fixed point in one second.

the distance between successive peaks in a wave.

Question 7

It takes light with a wavelength of 212 nm to break the NH bond in ammonia. What energy is required per photon to break this bond? What is the NH bond strength in terms of kJ per mole?

O 6.6 x 10⁻²² kJ/photon; 0.398 kJ/mol

0 6.6 x 10⁻²² kJ/photon; 398 kJ/mol

9.4 x 10⁻²² kJ/photon; 565,000 kJ/mol

9.4 x 10⁻²² kJ/photon; 565 kJ/mol

Question 8	1 pts
In 1 sec, a 60 W bulb emits 11 J of energy in the form of infrared radiation (heat) of wavelength 1850 nm. How mar photons of infrared radiation does the lamp generate in 1 sec?	ıy
◯ 1.02 x 10 ²⁰ photons	
◯ 1.04 x 10 ²⁹ photons	
◯ 1.10 x 10 ⁻¹⁹ photons	
O 6.63 x 10 ²³ photons	

1 pts

A photon has a frequency of 223 MHz. What is the energy of this photon?				
○ 1.48 x 10 ⁻²⁵ J				
◯ 1.48 x 10 ⁻³¹ J				
◯ 8.91 x 10 ⁻²⁸ J				
○ 8.91 x 10 ⁻²² J				

Question 10	1 pts
Carbon emits photons at 745 nm when exposed to blackbody radiation. How much energy would be obtained if 44 carbon were irradiated? Assume each carbon atom emits one photon.	4g of
○ 2.7 x 10 ⁻¹⁹ J	
○ 7.1 x 10 ⁶ J	
○ 9.1 x 10 ⁵ J	
◯ 5.9 x 10 ⁵ J	

Question 11	1 pts
A 200 nm photon has how many time the energy of a 700 nm photon?	
0.37	
0 4.2	
3.5	
0.29	

Question 12	1 pts

◯ 4.40 x 10 ⁻⁴³ J		
◯ 3.00 x 10 ⁻²⁵ J		
◯ 3.00 x 10 ⁻²² J		

Question 13	1 pts
Sodium vapor lamps, used for public lighting, emit yellow light of wavelength 570 nm. How much energy is emitted by excited sodium atom when it generates a photon?	' an
◯ 3.5 x 10 ⁻²⁸ J	
◯ 3.5 x 10 ⁻¹⁹ J	
◯ 2.8 x 10 ⁻¹⁹ J	
○ 2.8 x 10 ⁻²⁰ J	

Question 14	1 pts
Consider the sodium vapor lamps described in question 13. How much energy is emitted by 45.8 mg of sodium at emitting light at this wavelength? Assume each sodium atom emits one photon.	oms
◯ 2.0 x 10 ⁻³ J	
○ 4.2 x 10 ⁵ J	
○ 2.0 x 10 ²¹ J	
○ 420 J	

A particular metal has a work function of 1.05 eV. A light is shined onto this metal with a corresponding wavelength of 324 nm. What is the maximum velocity of the photoelectrons produced? (Hint: $1eV = 1.6022 \times 10^{-19}$ J, mass of an electron = 9.11×10^{-31} kg)

No photoelectrons are produced.		
◯ 1.16 x 10 ⁶ m/s		
\cap 1.35 x 10 ¹² m/s		

Question 16	1 pts
A particular metal has a work function of 3.05 eV. A light is shined onto this metal with a corresponding wavelent. What is the maximum velocity of the photoelectrons produced? (Hint: $1 \text{eV} = 1.6022 \times 10^{-19} \text{ J}$, mass of ar 9.11 x 10^{-31} kg)	ength of 524 າ electron =
○ 8.32 x 10 ¹¹ m/s	
O No photoelectrons are produced.	
○ 8.72 x 10 ⁸ m/s	
◯ 9.12 x 10 ⁵ m/s	

Question 17 1	l pts
Max Planck's theory averted the so called "UV Catastrophe" of classical mechanics. Which of the following best descri how Planck's theory avoided the "UV Catastrophe"?	bes
Radiation given off by blackbody radiators can only be emitted in quantized amounts.	
Radiation given off by blackbody radiators can be emitted in all types of radiation, not just UV radiation.	
Radiation emitted by blackbody radiators will reach UV energy levels only at extremely high temperatures.	
Eventually, blackbody radiators can cool to a temperature of absolute zero, resulting in its inability to release any more UV radiation.	

Question 18	1 pts
The de Broglie equation was important for a number of reasons, not least of which was that it demonstrated that	
O only quantum objects have wavelengths.	
all objects have a wavelength. However, in the case of quantum objects, these wavelengths are so small th they can be ignored.	at
O only macroscopic objects have wavelengths.	
all objects have a wavelength. However, in the case of macroscopic objects, these wavelengths are so smatter they can be ignored.	all that

Question 19	1 pts
An atom of which element, moving at 240 m/s, would possess a de Broglie wavelength of 1.40 x 10^{-11} m?	
◯ Cs	
◯ Sn	
O At	
O Mn	

Quiz saved at 9:44am	Submit Quiz