

1 H 1.008																	2 He 4.003
3 Li 6.941	4 Be 9.012											5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18
11 Na 22.99	12 Mg 24.31											13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.07	17 Cl 35.45	18 Ar 39.95
19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.87	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.69	29 Cu 63.55	30 Zn 65.38	31 Ga 69.72	32 Ge 72.64	33 As 74.92	34 Se 78.96	35 Br 79.90	36 Kr 83.80
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc (98)	44 Ru 101.07	45 Rh 102.91	46 Pd 106.42	47 Ag 107.87	48 Cd 112.41	49 In 114.82	50 Sn 118.71	51 Sb 121.76	52 Te 127.60	53 I 126.90	54 Xe 131.29
55 Cs 132.91	56 Ba 137.33	57 La 138.91	72 Hf 178.49	73 Ta 180.95	74 W 183.84	75 Re 186.21	76 Os 190.23	77 Ir 192.22	78 Pt 195.08	79 Au 196.97	80 Hg 200.59	81 Tl 204.38	82 Pb 207.20	83 Bi 208.98	84 Po (209)	85 At (210)	86 Rn (222)
87 Fr (223)	88 Ra (226)	89 Ac (227)	104 Rf (267)	105 Db (268)	106 Sg (269)	107 Bh (270)	108 Hs (270)	109 Mt (278)	110 Ds (281)	111 Rg (282)	112 Cn (285)	113 Nh (286)	114 Fl (289)	115 Mc (290)	116 Lv (293)	117 Ts (294)	118 Og (294)

58 Ce 140.12	59 Pr 140.91	60 Nd 144.24	61 Pm (145)	62 Sm 150.36	63 Eu 151.96	64 Gd 157.25	65 Tb 158.93	66 Dy 162.50	67 Ho 164.93	68 Er 167.26	69 Tm 168.93	70 Yb 173.04	71 Lu 174.97
90 Th 232.04	91 Pa 231.04	92 U 238.03	93 Np (237)	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (266)

Single Bond Energies

	H	C	N	O
H	436			
C	413	346		
N	391	305	163	
O	463	358	201	146

all values are kJ/mol

Multiple Bond Energies (kJ/mol)

C=C	602	C=N	615	C=O	799
C≡C	835	C≡N	887	C≡O	1072
N=N	418	N=O	607	C=S	577
N≡N	945	O=O	498		

water data

$C_{s,ice}$	= 2.09 J/g °C
$C_{s,water}$	= 4.184 J/g °C
$C_{s,steam}$	= 2.03 J/g °C
ρ_{water}	= 1.00 g/mL
ρ_{ice}	= 0.9167 g/mL
ΔH_{fus}	= 334 J/g
ΔH_{vap}	= 2260 J/g

This exam should have exactly 20 questions. Each question is equally weighted at 5 points each. Bubble in your answer choices on the bubble sheet provided. Your score is based on what you bubble on the bubblesheet and not what is circled on the exam.

1. A fuel is burned in a combustion reaction. You, safely standing in the surroundings, say, “wow that was hot!” This reaction is...

- a. exothermic
 - b. endothermic
-

2. A combustion reaction is performed in a bomb calorimeter. The temperature of the water rises from 25.1 °C to 26.9 °C. Which of the following is/are true regarding this process?

- I. the combustion reaction is exothermic
 - II. the combustion reaction is endothermic
 - III. the combustion reaction releases heat
 - IV. the heat flow of the calorimeter, q_{cal} , is positive
- a. I, II, and IV
 - b. I, III, and IV
 - c. I and IV
 - d. II and IV
 - e. I, II, III, and IV
-

3. Consider the following heat capacities for various substances:

$$\begin{array}{ll} C_{s,\text{lithium}} = 3.58 \frac{\text{J}}{\text{g} \cdot ^\circ\text{C}} & C_{s,\text{iron}} = 0.450 \frac{\text{J}}{\text{g} \cdot ^\circ\text{C}} \\ C_{s,\text{air}} = 1.012 \frac{\text{J}}{\text{g} \cdot ^\circ\text{C}} & C_{s,\text{helium}} = 5.193 \frac{\text{J}}{\text{g} \cdot ^\circ\text{C}} \\ C_{s,\text{mercury}} = 0.140 \frac{\text{J}}{\text{g} \cdot ^\circ\text{C}} & C_{s,\text{water}} = 4.184 \frac{\text{J}}{\text{g} \cdot ^\circ\text{C}} \end{array}$$

Which substance will have the greatest increase in temperature upon the addition of 10 kJ to the same mass of each substance?

- a. lithium
 - b. iron
 - c. mercury
 - d. air
 - e. water
 - f. helium
-

4. Determine which of the following processes are **endothermic**:

- I. a chemical reaction absorbs 44 kJ of heat
 - II. a cold glass of water freezes
 - III. silver jewelry is melted
- a. I and III
 - b. I and II
 - c. II only
 - d. I only
 - e. II and III
-

5. A standard gold bar held by central banks weighs approximately 12.4 kg. What is the minimum amount of energy needed to fully melt a metal bar that is already at its melting point? The heat of fusion for gold is 63.0 J/g.

- a. 781 kJ
 - b. 7.82 kJ
 - c. 197 kJ
 - d. 525 kJ
 - e. 5.25 kJ
 - f. 982 kJ
-

6. A 25 °C cup of water holds approximately 237 g. What is the final temperature of water if this cup of water loses 8.50 kJ?

- a. 16.4° C
 - b. 29.8 ° C
 - c. 12.5° C
 - d. 17.0° C
 - e. 8.50° C
-


7. Which of the following will take the most heat to raise 1 °C?

- a. 1.000 L water
 - b. 500 mL water
 - c. 250 mL water
 - d. These will all take the same amount of heat.
-

8. A 315 g sample of pure methanol liquid is sitting at 0 °C. What is the value of heat flow for the system if this liquid is cooled to its freezing point and then fully frozen? Assume the solid methanol is not cooled beyond its freezing point of -97.6° C. The specific heat capacity for methanol liquid is 2.14 J/g °C. The heat of fusion of methanol is 11.6 J/g and the heat of vaporization is 1100 J/g.

- a. -69.4 kJ
 - b. -65.8 kJ
 - c. -3.65 kJ
 - d. -3.47 kJ
 - e. +65.8 kJ
 - f. +74.0 kJ
 - g. +3.65 kJ
-


9. Consider the following balanced chemical equation:



If the heat of combustion for methanol is 726 kJ/mol, how much heat is being released in the balanced combustion reaction shown above?

- a. 1452 kJ
 - b. 726 kJ
 - c. 1089 kJ
 - d. 484 kJ
-

10. Consider the following balanced chemical equation:



How much heat is released when 32.7 g of butane are combusted? ΔH for the balanced reaction is equal to -4438.4 kJ/mol rxn. The molar mass of butane is 58.12 g/mol.

- a. 1249 kJ
 - b. 0.562 kJ
 - c. 4494 kJ
 - d. 2497 kJ
 - e. 76.47 kJ
 - f. 135.7 kJ
-


11. A simple dissolution reaction is performed in a coffee cup calorimeter. When 5.01 g NaCl is placed in 350 mL water, the temperature decreases by 0.227 °C. What is the heat of the dissolution reaction, q_{sys} ?

- a. +332 J
 - b. -332 J
 - c. +398 J
 - d. -398 J
 - e. +4.75 J
 - f. -4.75 J
-

12. When 0.432 g of a hydrocarbon fuel are combusted in a bomb calorimeter filled with 1.002 L water, a temperature increase of 0.991 °C is measured. What is the ΔH of the fuel in kJ/g? The heat capacity of the calorimeter hardware is equal to 2.04 kJ/ °C.

- a. -14.3 kJ/g
 - b. -6.18 kJ/g
 - c. +6.18 kJ/g
 - d. +3/18 kJ/g
 - e. -8.83 kJ/g
 - f. -1.79 kJ/g
-

13. Use bond energy data to determine the ΔH of the following balanced chemical equation:



- a. -93 kJ
 - b. +93 kJ
 - c. -990 kJ
 - d. +990 kJ
 - e. -620 kJ
 - f. 463 kJ
-

14. Which of the following is true regarding combustible fuels?

- a. The chemical reaction releases heat because the products are lower in energy than the reactants.
 - b. The chemical reaction absorbs heat because the products are lower in energy than the reactants.
 - c. The chemical reaction releases heat because the reactants are lower in energy than the products.
 - d. The chemical reaction absorbs heat because the reactants are lower in energy than the products.
 - e. The chemical reaction does not absorb or release heat because the products and reactants are equal in energy.
-

15. Consider the following heats of combustion for three fuel sources:

ΔH MTBE, 88.15 g/mol: 3,362.0 kJ/mol

ΔH Methane, 16.04 g/mol: 803.60 kJ/mol

ΔH Ethanol, 46.07 g/mol: 1234.7 kJ/mol

Which of these fuels is more efficient **per gram**?

- a. Methane
 - b. MTBE
 - c. They are equally efficient in kJ/g.
 - d. Ethanol
-

16. Asphalt is made from the largest carbon chains separated out in the crude oil refining process. In which portion of the distillation tower will you separate out these hydrocarbons?

- a. the bottom
 - b. the top
 - c. the middle
 - d. equally dispersed throughout the distillation tower
-

17. An ice cube is heated from $-15\text{ }^\circ\text{C}$ to steam at $115\text{ }^\circ\text{C}$. If you use a heating curve to solve for the total heat required for this process, how many unique heat calculations are necessary?

- a. 5
 - b. 6
 - c. 4
 - d. 3
 - e. 8
-

18. Select each phase change that requires heat flow into the system:

- I. freezing
- II. fusion
- III. vaporization
- IV. condensation

- a. I, II, and IV
 - b. I, III, and IV
 - c. I and IV
 - d. II and IV
 - e. I, II, III, and IV
 - f. II and III
-

19. Suppose you refined a large hydrocarbon sample and want to make more money out of your excessively large carbon chains. Which process will allow you to make a specific shorter carbon-based fuel?

- a. catalytic cracking
 - b. catalytic reforming
 - c. combustion
 - d. calorimetry
-

20. The process in which a hydrocarbon fuel can be further branched to increase fuel efficiency is called...

- a. cracking
- b. reforming
- c. combustion
- d. calorimetry

Remember to bubble in ALL your answers BEFORE time is called. Double check your name, utetid, and version number before you turn in your bubblesheet. You must keep your exam for future reference. Please do not lose it. We will not replace it.