4 pc	points $E = hv$ $c = \lambda v$
Long V	Microwave Infra-Red Vavelength, Low Energy requency UV X-Ray Gamma Ray Short Wavelength, High Energy High Frequency Increasing Energy (Ε), Frequency (ν) Shortening Wavelength (λ)
	the following radiation types from shortest to longest wavelength: K-Ray, Red light, Green light Green light, X-Ray, Red light, UV X-Ray, Green light, Red light, UV
	X-ray, UV, Green light, Red light Red light, Green light, UV, X-Ray points would you describe the most likely effect of visible light on a molecule?
	electron excitation molecular ionization molecular vibration molecular rotation
	would you describe the most likely effect of infra-red radiation on matter?
))) — _{4 n}	excitation vibration ionization
I DNA	A, generally considered a very stable organic polymer, is first damaged at which on of the electromagnetic spectrum? Visible light UV region Radio waves IR region
l Whic	ch of the following pairs the correct definition of frequency along with the correct shown in parenthesis? The number of wavelengths that pass a single point per second (s) The time it takes for a full wavelength to pass a single point (s ⁻¹) The time it takes for a full wavelength to pass a single point (s) The number of wavelengths that pass a single point per second (s ⁻¹)
	t is the frequency of yellow light with a wavelength of 580 nm? $2.39 \times 10^{-19} \text{ s}^{-1}$ $1.80 \times 10^{-7} \text{ s}^{-1}$ $5.17 \times 10^{14} \text{ s}^{-1}$ $5.17 \times 10^{5} \text{ s}^{-1}$
	th of the following equations directly solves for energy using wavelength? $E=h\lambda/\nu$ $E=h\lambda$ $E=hc/\lambda$ $E=hc/\lambda$
-M r ister	adio stations correspond to the frequency of the channel in MHz. If you want to a radio station that has a frequency equal to 1.015 x 10 ⁸ Hz, you should tune radio to 1015 10.15 101.5 1.015
I Micr wave	oints owaves, such as those used for radar and to heat food in a microwave oven, have elengths just greater than about 3 mm. What is the corresponding frequency of tion with a 13.4 mm wavelength?
0 0 0	$7.52 \times 10^{10} \text{Hz}$ $4.81 \times 10^{10} \text{Hz}$ $6.82 \times 10^{10} \text{Hz}$ $2.24 \times 10^{10} \text{Hz}$
l It tak ener	ces light with a wavelength of 212 nm to break the N-H bond in ammonia. What gy is required per photon to break this bond? What is the N-H bond strength in s of kJ per mole?
	9.38 x 10 ⁻²² kJ/photon; 565,000 kJ/mol 6.61 x 10 ⁻²² kJ/photon; 0.398 kJ/mol 9.38 x 10 ⁻²² kJ/photon; 565 kJ/mol 6.61 x 10 ⁻²² kJ/photon; 398 kJ/mol
	oints th of the following is the most energetic form of UV light? UV-B UV-A
_	UV-C UV-A, UV-B, and UV-C are equally energetic
	um vapor lamps, used for public lighting, emit yellow light of a wavelength of 57 much energy is emitted by an excited sodium atom when it generates a photon $2.8 \times 10^{-20} \text{J}$ $3.5 \times 10^{-19} \text{J}$ $2.8 \times 10^{-19} \text{J}$ $3.5 \times 10^{-28} \text{J}$
Cons s em	sider the sodium vapor lamps described in the previous question. How much end witted by 45.8 mg of sodium atoms emitting light at this wavelength? Assume early atom emits one photon. $4.2\times10^5\text{J}$ $2.0\times10^{21}\text{J}$ 420J $2.0\times10^{-3}\text{J}$
Whic	ch of the following statements are consistent with modern atomic theory? Multivers may apply. Electrons exist in discrete, quantifiable energy levels.
	An electron that has zero energy when it is closest to the nucleus The solutions to the Schrödinger Equation are wavefunctions that describe the energy and position of electrons in an atom. The vast majority of mass exists in the nucleus of an atom, but the radius of the nucleus is only about minuscule fraction of the overall atomic radius
	points ch of the following is a possible quantum number set for an electron in a 4d orbit $n=4,\ \ell=2,\ m_\ell=-1,\ m_{\rm S}=\frac{1}{2}$ $n=2,\ \ell=4,\ m_\ell=2,\ m_{\rm S}=-\frac{1}{2}$
0	$n = 4, \ \ell = 3, \ m_{\ell} = 3, \ m_{s} = \frac{1}{2}$ $n = 4, \ \ell = 3, \ m_{\ell} = 0, \ m_{s} = \frac{1}{2}$ points
An e	lectron is found in a 6f orbital. What is the value of the angular momentum quanter (\(\extrm{\ell} \) ? 6 3 1
)))	1 4 2 pints
	th of the following sets of quantum numbers is not possible? $n=2,\ l=0,\ m_l=0,\ m_s=-1/2$ $n=5,\ l=3,\ m_l=3,\ m_s=-1/2$ $n=3,\ l=2,\ m_l=1,\ m_s=1/2$
How	$n = 3$, $l = 4$, $m_l = -3$, $m_s = 1/2$ points many unique quantum number sets are possible for a 3p electron in an argon a
4 pc	points the following is not a possible quantum number set? $n = 3, \ \ell = 0, \ m_{\ell} = 0, \ m_{s} = \frac{1}{2}$
_	$n = 3, \ \ell = 0, \ m_{\ell} = 0, \ m_{S} = \frac{1}{2}$ $n = 3, \ \ell = 2, \ m_{\ell} = -3, \ m_{S} = -\frac{1}{2}$ $n = 4, \ \ell = 2, \ m_{\ell} = -1, \ m_{S} = -\frac{1}{2}$ $n = 2, \ \ell = 1, \ m_{\ell} = 0, \ m_{S} = \frac{1}{2}$
An e	points lectron orbital has a round, spherical shape (s-orbital). Its n value equals 3. What ible quantum number set for this electron orbital? n = 3, I = 1, mI = 0, ms = 1/2 n = 2, I = 3, mI = 0, ms = 1/2
	n = 5, l = 3, ml = 0, ms = 1/2 n = 3, l = 0, ml = 0, ms = 1/2
The G	electron configuration for the Mn atom is $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^7$ $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^5$ $1s^2 2s^2 2p^6 3s^2 3p^3$
	1s ² 2s ² 2p ⁶ 3s ² 3p ⁶ 4s ² 4p ⁵ points many total s electrons does P (atomic number 15) possess? 4
	 6 5 2
Wha	t is the electronic configuration of a selenium atom (Se)? $ [Ar]4s^23d^{10}4p^4 $ $ [Ar]4s^24d^{10}4p^4 $ $ [Kr]4s^24d^{10}4p^4 $ $ [Kr]4s^23d^{10}4p^4 $ $ [Kr]4s^23d^{10}4p^4 $ $ [Kr]4s^24p^{14}4d^{10}4p^4 $
Wha	t is the electron configuration of the chloride anion, C? $1s^22s^22p^63s^23p^4$ $1s^22s^22p^63s^23p^5$ $1s^22s^22p^63s^23p^6$ $1s^22s^22p^6$
3 pc	oints utral atom has a ground state electronic configuration designated 1s ² 2s ² 2p ² . Se tatement that best describes this atom.
\bigcirc	The atom has 2 unpaired electrons.

argon

magnesium

krypton

helium