version: 223

## Exam 2 - F22 - McCord - ch301

last name			f	first name				uteid				signature					
1																	18
1 H																	2 He
1.008	2											13	14	15	16	17	4.00
3	4	]										5	6	7	8	9	10
Li	Be											B	C	N	0	F	Ne
6.941 <b>11</b>	9.012 12	-										10.81 13	12.01 14	14.01 15	16.00 16	19.00 <b>17</b>	20.1 18
Na	М́д											AI	Si	P	S	CI	A
22.99	24.31	3	4	5	6	7	8	9	10	11	12	26.98	28.09	30.97	32.07	35.45	39.9
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
Κ	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	K
39.10	40.08	44.96	47.87	50.94	52.00	54.94	55.85	58.93	58.69	63.55	65.38	69.72	72.64	74.92	78.96	79.90	83.8
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rb	Sr	Y	Zr	Nb	Мо	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Те		Xe
85.47	87.62	88.91	91.22	92.91	95.94	(98)	101.07	102.91	106.42	107.87	112.41	114.82	118.71	121.76	127.60	126.90	131.
55	56	57	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	TI	Pb	Bi	Po	At	Rr
132.91	137.33	138.91	178.49	180.95	183.84	186.21	190.23	192.22	195.08	196.97	200.59	204.38	207.20	208.98	(209)	(210)	(22)
87	88	89	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118
Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Cn	Nh	FI	Mc	Lv	Ts	0
(223)	(226)	(227)	(267)	(268)	(269)	(270)	(270)	(278)	(281)	(282)	(285)	(286)	(289)	(290)	(293)	(294)	(294

58	59	60	61	62	63	64	65	66	67	68	69	70	71
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
140.12	140.91	144.24	(145)	150.36	151.96	157.25	158.93	162.50	164.93	167.26	168.93	173.04	174.97
90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
232.04	231.04	238.03	(237)	(244)	(243)	(247)	(247)	(251)	(252)	(257)	(258)	(259)	(266)

constants
R = 0.08206 L atm/mol K
$R=8.314~{\rm J/mol~K}$
$N_{\rm A} = 6.022 \times 10^{23} \ /{\rm mol}$
$h=6.626\times 10^{-34}~{\rm J\cdot s}$
$c=3.00\times 10^8~{\rm m/s}$
$F = 96485 \text{ C/mol e}^-$
$e = 1.602 \times 10^{-19} \text{ C}$
$m_{\rm e} = 9.11 \times 10^{-31} \text{ kg}$
Rydberg Constants
$\mathcal{R} = 2.18 \times 10^{-18} \text{ J}$
$\mathcal{R} = 3.29 \times 10^{15} \text{ s}^{-1}$
$\mathcal{R} = 1.097 \times 10^7 \text{ m}^{-1}$

## <u>conversions</u>

1  atm = 760  torr
$1~\mathrm{atm}=101325~\mathrm{Pa}$
1  atm = 1.01325  bar
1  atm = 14.7  psi
$1 \text{ bar} = 10^5 \text{ Pa}$
1  in = 2.54  cm
1  mi = 5280  ft
$1 \text{ Å} = 10^{-10} \text{ m}$
1  lb = 453.6  g
1  ton = 2000  lbs
1  tonne = 1000  kg
1  gal = 3.785  L
$1 \text{ gal} = 231 \text{ in}^3$
$1~{\rm fl}~{\rm oz}=29.57~{\rm mL}$

$$\label{eq:conversions} \begin{split} & \frac{\text{conversions}}{1 \ \text{cal} = 4.184 \ \text{J}} \\ & 1 \ \text{eV} = 1.602 \times 10^{-19} \ \text{J} \end{split}$$

1 kWh = 3600 J

## water data

$\overline{C_{\mathrm{s,ice}}} = 2.09 \mathrm{~J/g}$ °C
$C_{\rm s,water} = 4.184 \text{ J/g} \circ \text{C}$
$C_{\rm s,steam} = 2.03~\rm J/g~^\circ C$
$\rho_{\rm water} = 1.00~{\rm g/mL}$
$\rho_{\rm ice} = 0.9167~{\rm g/mL}$
$\rho_{\rm seawater} = 1.024~{\rm g/mL}$
$\Delta H_{\rm fus} = 334~{\rm J/g}$
$\Delta H_{\rm vap} = 2260~{\rm J/g}$
$K_{\rm w} = 1.0 \times 10^{-14}$

This exam should have exactly 25 questions. Each question is equally weighted at 4 points each. You will enter your answer choices on the virtual bubblehseet after you have finished. Your score is based on what you submit on the virtual bubblesheet and not what is circled on the exam.

1. Which of the following wavelengths corresponds to an emission band in the Balmer series?

•a.  $4.86 \times 10^{-7}$  m

b.  $5.56\times 10^{-9}~{\rm m}$ 

c.  $1.72\times 10^{-4}~{\rm m}$ 

- d. $9.53\times10^{-1}~{\rm m}$
- e. $6.88\times10^{-13}~{\rm m}$

**Explanation:**  $4.86 \times 10^{-7}$  m is the same as 486 nm. This is the only option within the visible spectrum, all of the emissions in the Balmer series occur in the visible region of electromagnetic radiation.

2. How many angular nodes would a theoretical 5g orbital have?

a. 3

b. 2

•c. 4

- d. 5
- e. 1

**Explanation:** A 5g orbital would have an angular momentum quantum number of  $\ell = 4$ , and thus would have 4 angular nodes.

**3.**What is the electron configuration of Bismuth (Bi)?

- a.  $[Xe]6s^{2}6p^{3}$
- b.  $[Rn]6p^{3}$
- c.  $[Rn]7s^25f^{14}6d^{10}7p^3$
- d.  $6s^24f^{14}5d^{10}6p^3$
- e.  $[Xe]6s^24f^{14}5d^{10}6p^3$
- f.  $[Xe]6s^25d^{10}6p^3$

**Explanation:** Electronic configurations can be abbreviated using the noble gases on the periodic table to respresent full valence shells. In the case of Bi, the closest previous noble gas is Xe. After the abbreviation, the electronic configuration is filled in by moving from left to right across the periodic table, taking care to not skip over the f-block elements.

4. The following species are isoelectronic. Select the atom or ion that will have the largest radius.

- a.  $Cl^-$
- b. Ar
- •c. S<sup>2-</sup>
- d.  $K^+$
- e.  $Ca^{2+}$

**Explanation:** Anions becomes larger with the addition of each electron due to electron repulsions in the valence shell. In this isoelectronic series, the sulfur ion has two extra electrons.

5. How many joules of energy are in one photon that has a wavelength of 166 nm?

•a.  $1.20 \times 10^{-18}$  J b.  $5.25 \times 10^{-19}$  J c.  $1.01 \times 10^{-40}$  J d.  $1.75 \times 10^{-18}$  J e.  $3.50 \times 10^{-19}$  J Explanation:  $E = hc/\lambda$  $= 6.626 \times 10^{-34} (3 \times 10^8)/166 \times 10^{-9} = 1.2 \times 10^{-18}$  J **6**. What is the electron configuration for the bromide ion (Br<sup>-</sup>)?

- a.  $[Ar]4s^24d^{10}4p^6$
- b.  $[Kr]4s^24d^{10}4p^4$
- c.  $[Ar]4s^24p^4$
- d.  $[Ar]4s^23d^{10}4p^4$
- e.  $[Ar]4s^23d^{10}4p^6$

**Explanation:** Begin at [Ar] and include only the electrons that fill after. This will include the 3d electrons and the one extra electron for the negative charge:  $[Ar]4s^23d^{10}4p^6$ 

7. What orbital property is the angular momentum quantum number  $(\ell)$  associated with?

- a. valence shell
- b. distance from nucleus
- c. electron spin
- •d. shape
- e. orientation in space

**Explanation:**  $\ell$  respresents the shape of the orbital, as determined by the angular nodes. The other answer choices are all associated with specific quantum numbers.

8. Which element/atom listed has the largest value for the effective nuclear charge  $(Z_{\text{eff}})$ ?

●a. Te

- b. Pb
- c. Sn
- d. Cs
- e. In

**Explanation:** The effective nuclear charge increases as you go across a row (period) of the periodic table. The number of inner shell electrons that *shield* the full nuclear charge stay constant across the row, but the number of protons is increasings, thus increasing  $Z_{\rm eff}$  and pulling the valence shell in tighter and smaller.

9. Which set of p orbitals below has been filled correctly? (they all have 4 electrons)

- a. <u>11 11</u> \_\_\_\_
- b. 11 1 1
- c.  $\underline{1} \downarrow \underline{1}$
- •d. <u>11 1 1</u>

**Explanation:** According to Hund's Rule, you fill singly before pairing begins. Also you must match the spin state when you do fill singly. So the first 3 electrons go in separate orbitals and they all have up spin (+1/2). The fourth electron pairs with the first and has down spin (-1/2).

10. An electron in the hydrogen atom is excited and makes the energy jump from n = 2 to n = 6. What wavelength of light corresponds to this energy jump?

- a.  $122~\mathrm{nm}$
- b. 656 nm
- •c. 410 nm
  - d. 434 nm
  - e. 486 nm

Explanation: Use the Rydberg equation:  $E = \mathcal{R} \left( \frac{1}{n^2} - \frac{1}{n^2} \right)$   $= 2.18 \times 10^{-18} (1/4 - 1/36) = 4.844 \times 10^{-19} \text{ J}$   $\lambda = \frac{hc}{E} = 6.626 \times 10^{-34} (3 \times 10^8) / 4.844 \times 10^{-19}$  $= 4.10 \times 10^{-7} \text{ m} = 410 \text{ nm}$ 

11. Household bleach contains sodium hypochlorite, what is the corresponding chemical formula?

- a. Na  $(ClO)_2$
- b.  $Na_2ClO$
- c.  $NaClO_2$
- d. NaClO<sub>3</sub>
- •e. NaClO

**Explanation:** Hypochlorite is the polyatomic ion  $ClO^-$ . To answer this question correctly, you also need to remember it has a -1 charge and that sodium always has a +1 charge as do all the alkali metals (group 1).

12. Television station KXAN in Austin is known as channel 36. That is in the UHF portion of communication frequencies. To be more specific, their video signal is on a carrier frequency of 603.25 MHz. Which of the following is the matching wavelength for this electromagnetic radiation?

a.  $20.8~\mathrm{cm}$ 

- b. 497  $\mu m$
- •c. 49.7 cm
- d.  $4.97~\mathrm{m}$
- e. 2.08 m

Explanation:  $\lambda = c/\nu$ = 3 × 10<sup>8</sup>/603.25 × 10<sup>6</sup> = 0.497 m = 49.7 cm

13. Which one of the atoms listed is paramagnetic?


●a. titanium, Ti

b. xenon, Xe

- c. cadmium, Cd
- d. mercury, Hg
- e. magnesium, Mg

**Explanation:** Ti has  $3d^2$  and those 2 electrons are not paired (Hund's Rule). This means that titanium is paramagnetic. The other choices all have matched (paired) sets of electrons completely filling a sublevel and making them all diamagnetic.

14. Below is a plot of sequential ionization energies. Which element is this?



**Explanation:** It is clear that the first 6 electrons follow an increasing trend and then there is a big jump on the 7th. This will only match well with sulfur and it's 6 valence electrons  $(3s^23p^4)$ . The 7th electron is a 2p electron and is much more difficult to remove to to its closer position to the nucleus. 15. Which quantum number set listed below could describe the highest energy valence electron of phosphorus?

a. n=2, ℓ=1
b. n=4, ℓ=0
c. n=3, ℓ=1
d. n=3, ℓ=0
e. n=3, ℓ=2

**Explanation:** The outermost filled shell (valence) is 3s and 3p. The 3p set is higher in energy than the 3s orbital. The 4s is higher than all of them, except it is empty - so it can't be right.

- 16. Which element has the largest atomic radius?
- a. S
- b. P
- c. Br
- •d. Sb
- e. Ar

**Explanation:** Atomic radii increase moving down a group(column) on the periodic table and decrease moving to the right across a period(row) on the periodic table. Moving down a row results in a larger increase in atomic radius than moving over 1 element to the left because moving down represents gaining an additional shell of electrons. Moving to the left just eases the contraction that occurs by having an additional electron in the valence shell interacting with the nucleus.

17. We are zapping various samples of metal with a laser ( $\lambda = 525$  nm). As we increase the work function of the metal, the velocity of the emitted electron...

- ●a. decreases
- b. increases
- c. it is impossible to know
- d. stays the same
- e. electrons aren't emitted, protons are

**Explanation:** The same wavelength of light is used in every case, so the energy of the incoming photon is constant. As the workfunction increases in energy, the kinetic energy of the emitted photoelectron decreases, resulting in slower velocities. 18. Which color of visible light corresponds to the highest frequency?

- a. orange
- b. red
- •c. blue
- d. green
- e. yellow

**Explanation:** Blue light is the high energy end of the visible light spectrum. Since higher energy postively correlates with higher frequency, blue light is the high frequency end of the visible light spectrum as well.

**19.** We conduct a photoelectric effect experiment using cesium and a light source of wavelength 540 nm. What is the velocity of the ejected electron from the surface of cesium? The work function ( $\Phi$ ) of cesium is 2.10 eV.

a. $6.96\times 10^{10}~{\rm m/s}$ 

- b.  $1.06\times 10^5~{\rm m/s}$
- c.  $3.11 \times 10^5$  m/s
- d.  $1.33 \times 10^{6} \text{ m/s}$
- e.  $2.64 \times 10^5$  m/s

## Explanation:

work function in J =  $2.1(1.602 \times 10^{-19}) = 3.3642 \times 10^{-19}$ incoming photon E =  $6.626 \times 10^{-34}(3 \times 10^8)/540 \times 10^{-9} = 3.6811 \times 10^{-19}$ difference is =  $3.1691 \times 10^{-20}$  J which is the kinetic energy of electron =  $1/2 \ mv^2$ 

 $v_{\text{electron}} = (2E_k/m)^{1/2}$ 

 $\left(2(3.1691\times 10^{-20})/9.11\times 10^{-31}\right)^{1/2} = 2.638\times 10^5~{\rm m/s}$ 

20.  $CCl_4$  was historically used as a flame retardant chemical in fire extinguishers. Given the chemical formula, what is the name of  $CCl_4$ ?

- •a. carbon tetrachloride
- b. carbon tetrachlorine
- c. carbon chloride
- d. carbon pentachloride
- e. chlorine tetracarbide

**Explanation:**  $CCl_4$  is a covalent compound. We start naming with carbon, as it is furthest left in the periodic table. Chlorine is next, but its suffix must be changed to -ide, it must also be given the correct prefix (tetra).

- •a. 5s
- b. 3s
- c. 6d
- d. 5d
- e. 4f
- f. 4s

**Explanation:** Radial nodes can be calculated using the equation:  $n-1-\ell$ . A 5s orbital would have 4 radial nodes, this is the highest of any of the options listed.

22. Suppose X stands for a particular element on the periodic table. Select the atom or ion that will have the largest radius.

- a.  $X^+$
- •b. X<sup>2-</sup>
- c.  $X^-$
- d.  $X^{2+}$
- e.  $\mathbf{X}$

**Explanation:** Anions becomes larger with the addition of each electron due to electron repulsions in the valence shell. The ion with the greatest negative charge will have the largest radius.

23. Which of the following choices correctly ranks the elements from the least electronegative to the most electronegative?

- a. F < Ca < N < Sr < Cs
- b.  $\mathrm{F} < \mathrm{N} < \mathrm{Ca} < \mathrm{Sr} < \mathrm{Cs}$
- c. Sr < N < F < Cs < Ca
- •d. Cs < Sr < Ca < N < F
- e. F < N < C<br/>s < Sr < Ca

**Explanation:** Electronegativity increases towards the top right corner of the periodic table, corresponding to the order: Cs < Sr < Ca < N < F.

page 6

24. Which of these ionic compounds has the lowest theoretical lattice energy?

a.  $CaCl_2$ 

b. CaO

 $c.\ MgCl_2$ 

d. NaBr

●e. KBr

**Explanation:** The lowest lattice energy will involve the lowest chargest and the largest radius. First, you can identify that KBr and NaBr have the lowest charges (+1/-1). Then KBr has the larger radius, making it weaker.

25. You meet Hank Hill, who sells propane and propane accessories. He asks if you, a chemistry student, can tell him the chemical formula of propane.

- a.  $\mathrm{C}_{4}\mathrm{H}_{10}$
- b.  $C_2H_6$
- c.  $C_3H_6$
- $d. \ C_4H_8$
- •e.  $C_3H_8$

**Explanation:** This is a pure nomeclature problem. Propane is the third alkane, corresponding to 3 carbons. The generic formula for all alkanes is  $C_nH_{2n+2}$ .

After you are finished and have all your answers circled, go to the front of the room and then use the QR code there to pull up the virtual answer page. Enter the appropriate info plus all your answers - click the SUBMIT button. Make sure you get the confirmation screen and show it to the TA or proctor. After that, turn in your exam and scratch paper. You're free to leave after that.



https://mccord.cm.utexas.edu/neon