version: 438 Exam 4 - F22 - McCord - ch301n

last na	me		f	irst na	me		u	teid		-	_	signa	ture				
1 1 H 1.008	2											13	14	15	16	17	18 2 He 4.003
3 Li 6.941	4 Be _{9.012}											5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne _{20.18}
11 Na 22.99	12 Mg _{24.31}	3	4	5	6	7	8	9	10	11	12	13 Al _{26.98}	14 Si _{28.09}	15 P 30.97	16 S 32.07	17 Cl 35.45	18 Ar 39.95
19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.87	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni _{58.69}	29 Cu 63.55	30 Zn 65.38	31 Ga _{69.72}	32 Ge	33 As _{74.92}	34 Se _{78.96}	35 Br 79.90	36 Kr 83.80
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc	44 Ru 101.07	45 Rh 102.91	46 Pd 106.42	47 Ag	48 Cd 112.41	49 In	50 Sn 118.71	51 Sb 121.76	52 Te	53 126.90	54 Xe 131.29
55 Cs 132.91	56 Ba	57 La	72 Hf 178.49	73 Ta 180.95	74 W 183.84	75 Re	76 Os 190.23	77 Ir 192.22	78 Pt 195.08	79 Au 196.97	80 Hg _{200.59}	81 TI 204.38	82 Pb 207.20	83 Bi _{208.98}	84 Po (209)	85 At (210)	86 Rn (222)
87 Fr (223)	88 Ra (226)	89 Ac (227)	104 Rf (267)	105 Db (268)	106 Sg (269)	107 Bh (270)	108 Hs (270)	109 Mt (278)	110 Ds (281)	111 Rg (282)	112 Cn (285)	113 Nh (286)	114 FI (289)	115 Mc (290)	116 Lv (293)	117 Ts (294)	118 Og (294)

58	59	60	61	62	63	64	65	66	67	68	69	70	71
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Но	Er	Tm	Yb	Lu
140.12	140.91	144.24	(145)	150.36	151.96	157.25	158.93	162.50	164.93	167.26	168.93	173.04	174.97
90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
232.04	231.04	238.03	(237)	(244)	(243)	(247)	(247)	(251)	(252)	(257)	(258)	(259)	(266)

constants

 $R=0.08206~\mathrm{L~atm/mol~K}$

R = 8.314 J/mol K

 $N_{\rm A} = 6.022 \times 10^{23} \ / {\rm mol}$

 $h=6.626\times 10^{-34}~\mathrm{J\cdot s}$

 $c = 3.00 \times 10^8 \text{ m/s}$

 $g = 9.81 \text{ m/s}^2$

conversions

1 atm = 760 torr

 $1~\mathrm{atm} = 101325~\mathrm{Pa}$

1 atm = 1.01325 bar

 $1 \text{ bar} = 10^5 \text{ Pa}$

 $^{\circ}F = ^{\circ}C(1.8) + 32$

 $K = {}^{\circ}C + 273.15$

conversions

1 in = 2.54 cm

1 ft = 12 in

1 yd = 3 ft

1 mi = 5280 ft

1 lb = 453.6 g

Single Bond Energies

	Н	С	N	О
Η	436			
\mathbf{C}	413	346		
Ν	391	305	163	
Ο	463	358	201	146

all values are kJ/mol

Multiple Bond Energies (kJ/mol)

C=C 602	C=N 615	C=O 799
$C \equiv C~835$	C≡ N 887	$C{\equiv}O~1072$
N=N 418	N = O 607	C=S 577
N≡N 945	O = O498	

water data

 $C_{\rm s,ice} = 2.09 \text{ J/g }^{\circ}\text{C}$

 $C_{\mathrm{s,water}} = 4.184 \mathrm{\ J/g\ ^{\circ}C}$

 $C_{\mathrm{s,steam}} = 2.03 \mathrm{\ J/g\ ^{\circ}C}$

 $\rho_{\rm water} = 1.00~{\rm g/mL}$

 $\rho_{\rm ice} = 0.9167~{\rm g/mL}$

 $\rho_{\rm seawater} = 1.024 \text{ g/mL}$

 $\Delta H_{\rm fus} = 334 \text{ J/g}$

 $\Delta H_{\rm vap} = 2260 \text{ J/g}$

 $K_{\rm w} = 1.0 \times 10^{-14}$

This exam should have exactly 25 questions. Each question is equally weighted at 4 points each. Bubble in your answer choices on the online bubblehseet provided. Your score is based on what you bubble on the bubblesheet and not what is circled on the exam.

- 1. A new fuel is found to have a heat of combustion of 37 kJ/g. The density of this fuel is 0.75 g/mL. How much heat is released when 20 mL of this fuel is burned?
- •a. 555 kJ
 - b. 612 kJ
 - c. 247 kJ
- d. 740 kJ
- e. 987 kJ

Explanation: (20 mL)(0.75 g/mL)(37 kJ/g) = 555 kJ

2. Consider the following molar masses and heats of combustion for three fuel sources:

 ΔH MTBE, 88.15 g/mol: 3,362.0 kJ/mol ΔH Methane, 16.04 g/mol: 803.60 kJ/mol

 ΔH Ethanol, 46.07 g/mol: 1234.7 kJ/mol

Which of these fuels is the least efficient **per gram** of fuel?

- a. Methane
- b. MTBE
- •c. Ethanol
- d. They are equally efficient in kJ/g.

Explanation: Convert the kJ/mol value into kJ/g for each and compare:

MTBE: 38 kJ/g = (3,362.0 kJ/mol)/(88.15 g/mol)Methane: 50.1 kJ/g = (803.60 kJ/mol)/(16.04 g/mol)

Ethanol: 27 kJ/g = (1234.7 kJ/mol)/(46.07 g/mol)

- 3. A reaction performed in a coffee cup calorimeter makes the temperature of the water increase. Which of the following is/are true regarding this process?
- I. the reaction is exothermic
- II. the reaction is endothermic
- III. the reaction absorbs heat
- IV. the value of $q_{\rm cal}$ is positive
- a. I, II, III, and IV
- •b. I and IV
 - c. I, III, and IV
 - d. II and IV
 - e. II, III, and IV

Explanation: For this process, the $q_{\rm cal}$ is positive (it absorbed heat which is why the temperature increased. Flip the sign to get into the perspective of the reaction system. That means that the reaction is exothermic and releases heat. Only I and IV are correct.

- 4. Use bond energy data (front page of exam) to determine the heat of combustion of acetylene, C₂H₂.
- a. -1449 kJ/mol
- b. -1705 kJ/mol
- c. -1300 kJ/mol
- d. -312 kJ/mol
- ●e. -1216 kJ/mol

Explanation: balance rxn: $C_2H_2 + 2.5O_2 \longrightarrow 2CO_2 + H_2O$

Reactant side: 2 C-H single bonds, 1 C \equiv C triple bond, 2.5 O=O double bonds. Product side: 4 C=O double bonds, 2 H-O single bonds.

Reactants = 2(413) + 1(835) + 2.5(498) = 2906 kJ

Products = 2(463) + 4(799) = 4122 kJ

Reactants - Products = -1216 kJ

- 5. A sample container holding 125 grams of water is heated from 24 °C to steam at 105 °. Calculate the heat necessary for this process. (answer to nearest whole kJ)
- •a. 324 kJ
- b. 83 kJ
- c. 41 kJ
- d. 298 kJ
- e. 361 kJ

Explanation: Three calculations are necessary here:

$$q_1 = (125 \,\mathrm{g})(4.184 \,\mathrm{J/g})(76^{\circ} \,\mathrm{C}) = 39748 \,\mathrm{J}$$

$$q_2 = (125 \text{ g})(2260 \text{ J/g}) = 282500 \text{ J}$$

$$q_3 = (125 \,\mathrm{g})(2.03 \,\mathrm{J/g})(5^{\circ} \,\mathrm{C}) = 1268.75 \,\mathrm{J}$$

$$q_{\rm total} = 39748 \; {\rm J} + 282500 \; {\rm J} + 1268.75 \; {\rm J} \approx 324 \; {\rm kJ}$$

- **6.** An *open* thermodynamic system allows for the transfer of what across a system boundary?
- a. Both matter and energy
- b. Energy only
- c. Neither matter nor energy
- d. Matter only

Explanation: An open system allows both energy and matter to flow in and out of the system boundary.

- 7. A calorimeter is only useful for measuring the energy content of exothermic reactions, not endothermic ones.
- •a. false
 - b. true

Explanation: Works well for any kind of reaction, endothermic or exothermic.

- 8. When a petroleum engineer proclaims "it's time to get crackin'!" they really mean that ...
- a. it's time to make the hydrocarbons more branched
- b. it's time to make the meth
- c. it's time to break larger hydrocarbons into smaller ones
- d. it's time to build bigger hydrocarbons from smaller ones

Explanation: cracking is breaking large hydrocarbons into smaller ones about half their original size

- 9. This is the corrected version of this question. All students received credit for this due to two typos. A beaker contains 288 g of methanol at 28.0 °C. What is the final temperature of methanol if it is cooled and loses 6.41 kJ of heat? The specific heat of methanol is 2.53 J/g°C.
- a. 8.50° C
- b. 21.8.° C
- c. 16.4° C
- d. 29.8 ° C
- ●e. 19.2° C

Explanation: Use the formula: $q = mC_s\Delta T$

$$-6410 \text{ J} = (288 \text{ g})(2.53 \text{ J/g}^{\circ} \text{ C})(T_f - 28^{\circ} \text{ C})$$

$$T_f = 19.2 \, ^{\circ}C$$

- 10. I want my room temperature beverage to be cold, so I put it in the refrigerator for an hour. Then pull it out and drink it yes, it is much colder now. I, being smart chemistry guy, know that if the beverage is the system and it "got cold" in the refrigerator, then the process must be exothermic.
- a. false
- •b. true

Explanation: Yes, it is true because heat was released from the beverage into the surroundings (the refrigerator) which means the process is exothermic.

- 11. Methane, ethane, propane, and butane are all examples of refinery gases. Based on the fact that these gases are the shortest carbon chains, where would you find these in a distillation tower?
- a. the bottom
- •b. the top
 - c. equally dispersed throughout the distillation tower
 - d. the middle

Explanation: Light chains remain at the higher portions of the distillation tower.

- 12. Which of the following processes would have a positive value for heat flow?
- a. deposition
- b. freezing
- c. condensation
- •d. sublimation

Explanation: Sublimation is an endothermic process, converting a solid directly to a gas. Such a process requires heat input which means a positive q and ΔH for the process. The other three are all exothermic.

- 13. The standard heat of combustion of ethanol, C_2H_5OH , is 1372 kJ/mol ethanol. How much heat (in kJ) would be liberated by completely burning a 20.0 g sample?
- a.686 kJ
- b. 469 kJ
- c. 519 kJ
- ●d. 597 kJ
- e. 715 kJ

Explanation: 20.0 g ethanol / 46 g/mol = 0.43 moles of ethanol. $1372 \text{ kJ/mol} \times 0.43 \text{ mol} = 597 \text{ kJ}.$

- 14. When 0.328 g of a hydrocarbon fuel are combusted in a bomb calorimeter filled with 925 mL water, a temperature increase of 2.15 °C is measured. What is the ΔH of the fuel in kJ/g? The heat capacity of the calorimeter hardware is equal to 1550 J/°C.
- •a. -35.5 kJ/g
 - b. -45.2 kJ/g
- c. +55.1 kJ/g
- d. +48.4 kJ/g
- e. -52.9 kJ/g
- f. -39.7 kJ/g

Explanation: bomb calorimetry: $\Delta H \approx -q_{\rm cal}$

$$q_{\rm cal} = mC\Delta T + C\Delta T$$

11653 = (925)(4.184)(2.15) + (1550)(2.15)

Flip the sign to get into the system: -11653 J

Convert to kJ and divide by mass:

-11.653 kJ/0.328 g = -35.5 kJ/g

- 15. Which of the following is true regarding combustible fuels?
- a. The chemical reaction absorbs heat because the products are lower in energy than the reactants.
- b. The chemical reaction absorbs heat because the reactants are lower in energy than the products.
- c. The chemical reaction does not absorb or release heat because the products and reactants are equal in energy.
- •d. The chemical reaction releases heat because the products are lower in energy than the reactants.
- e. The chemical reaction releases heat because the reactants are lower in energy than the products.

Explanation: The driving force behind burning a fuel is the fact that the chemical reaction releases heat because the products are lower in energy than the reactants.

- 16. The first law of thermodynamics is all about the
- a. conversion of mass to energy
- b. generation of antimatter
- c. dispersal of energy
- •d. conservation of energy

Explanation: conservation of energy

17. Which process of crude oil refining would most efficiently convert this linear molecule of n-octane to a branched molecule called 2,5-dimethylhexane?

$$\longrightarrow$$

- a. catalytic cracking
- b. fractional distillation
- •c. catalytic reforming
 - d. thermal cracking

Explanation: Catalytic reforming is used to break down linear chain hydrocarbons and remake them into highly branched ones.

- 18. Which of the following is an exothermic process?
- •a. Freezing water
 - b. Melting ice
 - c. Heating water
 - d. Boiling water

Explanation: Heat is released when water is frozen to a solid. The other three absorb heat, which is endothermic.

19. Consider the following heat capacities (all in J/g°C) for various substances:

 $C_{\rm s,chromium} = 0.449$

 $C_{\text{s.titanium}} = 0.523$

 $C_{\rm s, water} = 4.184$

 $C_{\rm s.air} = 1.012$

Which substance will have the greatest increase in temperature upon the addition of 275 joules of heat to the same mass of each substance?

- a. air
- b. water
- c. titanium
- d. chromium

Explanation: The greatest temperature change will be observed in the substance with the lowest specific heat capacity. This is titanium.

- 20. Which of the following is the least efficient fuel based on its combustion enthalpy per gram?
- a. hydrogen
- b. methane
- c. propane
- •d. wood
- e. ethanol

Explanation: Wood can only produce about 14 kJ of energy per gram, which is why its the least efficient.

- 21. The second law of thermodynamics states that the energy of the universe is always increasing.
- a. true
- •b. false

Explanation: No. The entropy of the universe is increasing due to spontaneous changes. First law says the energy of the universe is constant.

- 22. Which process can turn a compound that is found lower on a distillation tower into a a more desirable compound that would be found higher on the distillation tower?
- a. catalytic reforming
- b. fracking
- c. combustion
- d. catalytic cracking
- e. calorimetry

Explanation: Catalytic cracking can provide shorter carbon chains from longer carbon chains in a specific, efficient manner.

- 23. When a hydrocarbon for a combustion engine is more branched $___$.
- •a. it burns smoother and will have a higher octane
 - b. it will have a much higher energy of combustion
 - c. it will fail to combust
 - d. it will more readily pre-ignite which causes knocking

Explanation: more branching is desired (reforming) which leads to better burning fuels and higher octane ratings

- **24.** A simple dissolution reaction is performed in a coffee cup calorimeter. When 5.12 g of a soluble ionic salt is dissolved into 180 mL water, the temperature decreases by 9.44 °C. What is the value of ΔH for this dissolution process?
- a. +5.81 kJ
- b. +3.92 kJ
- c. -5.81 kJ
- d. -7.11 kJ
- e. -3.92 kJ
- ●f. +7.11 kJ

Explanation: Solve for q_{cal} and then flip the sign for q_{sys} .

- $q = (180 \,\mathrm{g})(4.184 \,\mathrm{J/g}\,^{\circ}\mathrm{C})(-9.44\,^{\circ}\mathrm{C})$
- q = -7109 J = -7.11 kJ

Flip the sign to get back to the system to get +7.11 kJ.

- 25. Fill in the blanks: Breaking bonds is an _____ process, and making bonds is an _____ one. Thus, the ΔH values for breaking and making bonds respectively, are ____ and ____ .
- •a. endothermic, exothermic, positive, negative
- b. endothermic, exothermic, negative, positive
- c. exothermic, endothermic, positive, negative
- d. exothermic, endothermic, negative, positive

Explanation: Breaking bonds requires energy input, so it is an endothermic process, which has a positive ΔH . The opposite is true for making bonds.

After you are finished and have all your answers circled, go to the front of the room and then use the QR code there to pull up the virtual answer page. Enter the appropriate info plus all your answers - click the SUBMIT button. Make sure you get the confirmation screen and show it to the TA or proctor. After that, turn in your exam and scratch paper. You're free to leave after that.



https://mccord.cm.utexas.edu/helium

* * * Remember your Version Number * * *