

## HW04 - Chemical Equilibria 2

### Question 1

2 pts


Calculate the equilibrium constant at 25°C for a reaction for which  $\Delta G^\circ = -4.22$  kcal/mol.

- 620.254
- 2481.02
- 1240.51
- 1240.51

### Question 2

2 pts

The reaction



has an equilibrium constant of  $3.7 \times 10^{-3}$ . Consider a reaction mixture with:

$$[A] = 2.0 \times 10^{-2} \text{ M}$$

$$[B] = 1.7 \times 10^{-4} \text{ M}$$

$$[C] = 2.4 \times 10^{-6} \text{ M}$$

$$[D] = 3.5 \times 10^{-3} \text{ M}$$


Which of the following statements is definitely true?

- The reverse reaction will occur to a greater extent than the forward reaction until equilibrium is established.
- The forward reaction will occur to a greater extent than the reverse reaction until equilibrium is established.
- No conclusions about the system can be made without additional information.
- The system is at equilibrium.

### Question 3

2 pts

The reaction




has an equilibrium constant ( $K_c$ ) of  $4.0 \times 10^8$  at 25°C. What will eventually happen if 44.0 moles of  $NH_3$ , 0.452 moles of  $N_2$ , and 0.108 moles of  $H_2$  are put in a 10.0 L container at 25°

- It is impossible to know what will happen unless we know what the equilibrium constant is at 298 K.
- More  $N_2$  and  $H_2$  will be formed.
- More  $NH_3$  will be formed.
- Nothing. The system is at equilibrium.

### Question 4

2 pts

Consider the reaction:



If the initial concentration of  $Ni(CO)_4(g)$  is 1.0 M and  $x$  is the equilibrium concentration of  $CO(g)$ , what is the correct equilibrium relation?

- $K_c = \frac{256x^4}{(1.0 - 4x)}$
- $K_c = \frac{x^4}{(1.0 - \frac{x}{4})}$
- $K_c = \frac{x^5}{(1.0 - \frac{x}{4})}$
- $K_c = \frac{4x}{(1.0 - 4x)}$

### Question 5

2 pts

Suppose the reaction




has an equilibrium constant  $K_c = 49$  and the initial concentrations of  $H_2$  and  $I_2$  is 0.5 M and of  $HI$  is 0.0M. Which of the following is the correct value for the final concentration of  $HI(g)$ ?

- 0.778 M
- 0.219 M
- 0.250 M
- 0.599 M

### Question 6

2 pts

The system



is at equilibrium at a fixed temperature with a partial pressure of  $H_2$  of 0.200 atm, a partial pressure of  $I_2$  of 0.200 atm, and a partial pressure of  $HI$  of 0.100 atm. An additional 0.26 atm pressure of  $HI$  is admitted to the container, and it is allowed to come to equilibrium again. What is the new partial pressure of  $HI$ ?

- 0.104 atm
- 0.464 atm
- 0.152 atm
- 0.360 atm

### Question 7

2 pts

At 990°C,  $K_c = 1.6$  for the reaction




How many moles of  $H_2O(g)$  are present in an equilibrium mixture resulting from the addition of 1.00 mole of  $H_2$ , 2.00 moles of  $CO_2$ , 0.75 moles of  $H_2O$ , and 1.00 mole of  $CO$  to a 5.00 liter container at 990°C?

- 1.1 mol
- 1.0 mol
- 0.60 mol
- 1.7 mol

### Question 8

2 pts

What happens to the concentration of  $NO(g)$  when the total pressure on the reaction below is increased (by compression) when it is at equilibrium?




- it remains the same
- it decreases
- it increases
- it is impossible to tell

### Question 9

2 pts

Consider the following reaction:




where  $\Delta H_{rxn} = -198$  kJ. The amount of  $SO_2(g)$  at equilibrium increases when...

- $SO_3$  is removed.
- the volume is increased.
- more oxygen is added.
- the temperature is decreased.

### Question 10

2 pts

Suppose the reaction mixture



is at equilibrium at a given temperature and pressure. The pressure is then increased at constant temperature by compressing the reaction mixture, and the mixture is then allowed to reestablish equilibrium. At the new equilibrium...

- there is less ammonia present than there was originally.
- there is the same amount of ammonia present as there was originally.
- there is more ammonia present than there was originally.
- the nitrogen is used up completely.

### Question 11

2 pts

Consider the system:




at equilibrium at 25°C. If this is an exothermic reaction and the temperature was raised, would the equilibrium be shifted to produce more  $N_2O_5$  or more  $N_2O_4$ ?

- more  $N_2O_5$
- there would be no change
- more  $N_2O_4$
- it is impossible to tell

### Question 12

2 pts

The system




is at equilibrium at some temperature. At equilibrium, a 4.00L vessel contains 1.00 mole  $CO_2$ , 1.00 mole  $H_2$ , 2.40 moles  $H_2O$ , and 2.40 moles  $CO$ . How many moles of  $CO_2$  must be added to this system to bring the equilibrium  $CO$  concentration to 0.669 mol/L?

- 0.429 moles
- 0.498 moles
- 0.069 moles
- 0.993 moles

### Question 13

1 pts

The figure below represents a reaction at 298 K.



Based on the figure, which of the following statements (if any) are FALSE?

- At point D, the reaction will move toward the reactants to get to equilibrium.
- For this reaction,  $\Delta G^\circ$  is negative.
- At point C, the system is at equilibrium.
- None of the other statements are false.
- At point B,  $Q < K$ .

### Question 14

1 pts

Given the hypothetical reaction:



Predict what will happen when 1.0 mol  $Y$  is placed into an evacuated container.

- $Q$  will increase until  $Q = K$ .
- Nothing. The products are already formed, so no reaction occurs.
- $Q$  will decrease until  $Q = K$ .
- $\Delta G^\circ$  will decrease until  $\Delta G^\circ = 0$ .

### Question 15

2 pts

Consider the reaction:



Which of the following is a possible value of  $K$  for this reaction?

- 0.56
- $10^{70}$
- $10^{-70}$
- 0.56

### Question 16

2 pts

The equilibrium constant  $K$  for the synthesis of ammonia is  $6.8 \times 10^5$  at 298 K. What will  $K$  be for the reaction at 375 K?



- 326
- $1.42 \times 10^9$
- $6.75 \times 10^5$
- $6.85 \times 10^5$