

This print-out should have 50 questions. Multiple-choice questions may continue on the next column or page – find all choices before answering.


001 10.0 points

What is the dominant species in solution at the equivalence point of a weak base-strong acid titration?

1. Weak acid
2. Strong base
3. Neutral salt
4. Weak base
5. Equal parts weak acid and weak base
6. Strong acid

002 10.0 points

A and B are mildly volatile solvents. A mixture is made by combining 2 moles of A with 3 moles of B. Interpret the diagram below to determine the vapor pressure of this mixture.



1. 90 Torr
2. 110 Torr
3. 130 Torr
4. 150 Torr
5. 140 Torr
6. 70 Torr
7. 120 Torr

8. 80 Torr

9. 100 Torr


003 10.0 points

A reaction has a negative change in entropy. This reaction can only be spontaneous if...

1. heat is absorbed at a sufficiently high temperature
2. heat is released at a sufficiently low temperature
3. heat is released at any temperature
4. heat is absorbed by the system at any temperature
5. None of these choices are correct because a reaction with a negative change in entropy can never be spontaneous

004 10.0 points

Consider the following overall reaction:



Using the overall reaction, determine the rate law for the following mechanism:



1. Rate = $k' [Z] [X]$
2. Rate = $k' [A_2] [Z] [X]$
3. Rate = $k' [A_2]^2$
4. Rate = $k' [A_2]^2 [X]$
5. Rate = $k' [A_2] [X]$
6. Rate = $k' [Z][Y]$

005 10.0 points

Using an electroplating system operating at

7.35 amps, it take 1.50 hours to plate out 5.00 grams of an unknown metal from its molten chloride salt, MCl_2 . Identify the metal M.

1. Mg
2. Cd
3. Cu
4. Fe
5. Zn

006 10.0 points

A 19.7 g sample of an unknown salt (formula = MX_2) is dissolved in 249.4 mL water. The boiling point of water in this solution is 100.657°C . What is the molecular weight of the unknown salt?

1. 185 g/mol
2. 61.6 g/mol
3. 46.1 g/mol
4. 129.8 g/mol
5. 55.4 g/mol

007 10.0 points

Barium fluoride (BaF_2) is most soluble in which of the following solutions?

1. 0.50 M NaF
2. 0.005 M $Ba(OH)_2$
3. 0.18 M NaF
4. The molar solubility of barium fluoride is the same in each of these solutions
5. 0.15 M $Ba(OH)_2$

008 10.0 points

Consider the following substances: acetic acid (CH_3COOH), propane (C_3H_8), and acetone

(CH_3COCH_3). The boiling points (in no particular order) are -42°C , 56°C , and 118°C . The vapor pressures (in no particular order) are 225 Torr, 15 Torr, and 6400 Torr. What is the boiling point and vapor pressure for acetic acid?

1. 118°C , 15 Torr
2. 56°C , 225 Torr
3. -42°C , 15 Torr
4. -42°C , 6400 Torr
5. 118°C , 6400 Torr


009 10.0 points

What is the ratio of potassium acetate to acetic acid necessary to make a buffer with a pH equal to 5.12?

1. 2.4
2. 3.1
3. 0.38
4. 0.42
5. 2.9
6. 9.86

010 10.0 points

Consider the following reaction:




Oxygen is being consumed at a rate equal to 1.24 M/s. What is the initial rate at which carbon dioxide is forming?

1. 1.24 M/s
2. 4.96 M/s
3. 0.709 M/s
4. 8.68 M/s

5. 2.17 M/s

011 10.0 points

Identify the missing isotope in the nuclear reaction.

1. ${}_{86}^{222}\text{Rn}$ 2. ${}_{86}^{226}\text{Rn}$ 3. ${}_{86}^{230}\text{Th}$ 4. ${}_{90}^{230}\text{Th}$ 5. ${}_{90}^{222}\text{Rn}$

012 10.0 pointsWhat is the pH after 250 mL of 0.25 M HNO_3 is added to 350 mL of 0.50 M CH_3NH_2 (methylamine) ?

1. 3.36

2. 10.90

3. 10.64

4. 6.21

5. 2.29


6. 10.38

7. 3.62

8. 5.05

013 10.0 points

0.834 atm A, 0.565 atm B, and 1.24 atm C are placed into a container to run the following reaction:

At equilibrium, 0.435 atm C remains. What is K_p for this reaction?

1. 18.8

2. 0.0526

3. 0.107

4. 0.0230

5. 8.12


014 10.0 points

A weak acid, HA, ionizes 7.65% at a 0.250 M concentration. What is the hydroxide ion concentration in this solution?

1. 1.00×10^{-14} M2. 1.91×10^{-2} M3. 5.23×10^{-13} M4. 1.31×10^{-13} M5. 1.91×10^{-16} M


015 10.0 points




The following diagram shows a solution on the left (dark shade) and just the solvent on the right (light shade) separated by a semipermeable membrane.



Which diagram best represents the final state of this system after equilibrium is achieved?


1.





016 (part 1 of 2) 10.0 points

You run an experiment to determine the initial rates of the following generic reaction at various starting conditions:



	[A] M	[X ₂] M	initial rate M · s ⁻¹
Trial 1	0.60	1.56	2.00 × 10 ⁻³
Trial 2	0.60	3.12	8.00 × 10 ⁻³
Trial 3	1.20	1.56	4.00 × 10 ⁻³
Trial 4	0.90	2.40	7.10 × 10 ⁻³

What is the correct rate law for the reaction?

1. Rate = (3.84 × 10⁻³)[A]⁻¹[X₂]²
2. Rate = (1.40)[A]
3. Rate = (1.37 × 10⁻³)[A][X₂]
4. Rate = (3.84 × 10⁻³)[A][X₂]²
5. Rate = (1.37 × 10⁻³)[A][X₂]²


017 (part 2 of 2) 10.0 points

What are the units of the rate constant in the previous question?

1. $\frac{1}{M^3 \cdot s}$
2. $\frac{1}{M \cdot s}$
3. $\frac{M}{s}$
4. $\frac{1}{M^2 \cdot s}$
5. $\frac{1}{M^4 \cdot s}$

018 (part 1 of 2) 10.0 points

What is the shorthand notation for the following electrochemical cell?



1. Cr²⁺ | Cr³⁺ || Co²⁺ | Co
2. Cr²⁺ , Cr³⁺ || Co²⁺ | Co
3. Pt | Cr²⁺ , Cr³⁺ || Co²⁺ | Co
4. Co | Co²⁺ || Cr²⁺ , Cr³⁺ | Pt
5. Co²⁺ | Co || Cr²⁺ , Cr³⁺


019 (part 2 of 2) 10.0 points

What is the oxidizing agent in the previous problem?

1. Co²⁺
2. Pt
3. Cr³⁺
4. Co
5. Cr²⁺

020 (part 1 of 2) 10.0 points

Use the following phase diagram for the next two questions.



What is the normal melting point for this substance? Note: the vertical axis is logarithmic in scale.

1. 0°C
2. 150°C
3. 75°C
4. 120°C
5. 20°C
6. 45°C

021 (part 2 of 2) 10.0 points

A sample of this substance is held at 0.1 atm and -50°C . The sample is pressurized to 3 atm and then heated to 250°C . In total, what phase transitions occurred?

1. melting and boiling
2. sublimation and condensation
3. melting and condensation
4. sublimation only
5. melting and freezing

022 10.0 points

Use half-reactions from the standard reduction table to calculate the K_{sp} for $\text{Zn}(\text{IO}_3)_2$.

1. 3.9×10^{-6}
2. 4.8×10^{-12}
3. 1.7×10^{-26}
4. 6.8×10^{-32}
5. 7.3×10^{-19}

023 10.0 points

^{123}I is a radioactive isotope ($t_{1/2} = 13.22$ hours) useful for clinical imaging. How long will it take for a dose to diminish to 18.7% of its original value?

1. 34.2 hours
2. 557 hours
3. 38.7 hours
4. 87.9 hours
5. 70.7 hours
6. 29.7 hours
7. 22.2 hours
8. 32.0 hours

024 10.0 points


Which of the following salts is the most soluble in pure water?

1. CuBr $K_{\text{sp}} = 6.3 \times 10^{-9}$
2. CaF_2 $K_{\text{sp}} = 3.5 \times 10^{-11}$
3. BaSO_4 $K_{\text{sp}} = 1.1 \times 10^{-10}$
4. All of these salts have the same solubility

025 10.0 points

Write the equilibrium constant for the follow-

ing reaction.



1. $K_p = \frac{P_{\text{HBr}}^2}{P_{\text{H}_2}}$

2. $K_p = \frac{P_{\text{HBr}}^2}{P_{\text{H}_2}[\text{Br}_2]}$

3. $K_p = \frac{P_{\text{H}_2}}{P_{\text{HBr}}^2}$

4. $K_p = \frac{P_{\text{HBr}}}{P_{\text{H}_2}}$

5. $K_p = \frac{P_{\text{HBr}}^2}{P_{\text{H}_2}P_{\text{Br}_2}}$

026 10.0 points

A sample of 44.1 g of *para*-dichlorobenzene ($\text{C}_6\text{H}_4\text{Cl}_2$, 147.0 g/mol) is dissolved into 350 mL of hexane (C_6H_{14} , 86.18 g/mol, density 0.661 g/mL). What is the molality of this solution?

1. 1.30 *m*

2. 0.101 *m*

3. 0.857 *m*

4. 1.17 *m*

5. 0.567 *m*

027 10.0 points

Calculate the number of moles of oxygen that will dissolve in 45 L of water at 20°C if the partial pressure of oxygen is 0.21 atm. The Henry's Law constant for oxygen in water at 20°C is 0.0013 M/atm.

1. 0.0062 mol

2. 0.00027 mol

3. 0.0013 mol

4. 0.28 mol

5. 0.012 mol

028 10.0 points

What is the pH of a 0.12 M solution of hydroxylammonium bromide (NH_3OHBr)?

1. 4.44

2. 3.24

3. 3.33

4. 3.61

5. 9.56


6. 3.48

7. 3.72

8. 6.04

029 10.0 points

The following reaction occurs in acidic conditions. What is the coefficient of water in the overall balanced equation? Is it a reactant or a product?



1. 3; reactant

2. 7; reactant

3. 2; product


4. 4; reactant

5. 2; reactant

6. 3; product

030 10.0 points


What is the potential for the following cell?



1. 2.37 V
2. 1.88 V
3. 2.02 V
4. 1.95 V
5. 1.81 V
6. 2.09 V

031 (part 1 of 2) 10.0 points

Consider the following reaction for the next two questions:



Beginning at equilibrium, which of the following will result in a shift toward the right of this reaction?

1. Adding an inert gas at constant volume
2. Adding an inert gas at constant pressure
3. Adding CH_3OH gas
4. Removing CO gas
5. Reducing the volume of the container

032 (part 2 of 2) 10.0 points


If this reaction is exothermic, lowering the temperature will cause the reaction to...

1. remain at equilibrium
2. shift right due to a larger K value
3. shift left due to a smaller K value
4. shift left due to a smaller Q value
5. shift right due to a decreased Q value

033 (part 1 of 2) 10.0 points

The following two questions refer to this diagram for a voltaic cell. Neither of the two

electrodes are an inert electrode.



Where would you find the species that is being oxidized?

1. D
2. C
3. B
4. A
5. E

034 (part 2 of 2) 10.0 points



If the half-reaction for the anode involves Fe^{2+} and Fe , which of these redox pairs could be in the cell on the right?

1. H^+ and H_2
2. None of these can give a voltaic cell
3. Cr^{3+} and Cr
4. Sn^{2+} and Sn
5. Mn^{2+} and Mn


035 10.0 points

$^{99}_{42}\text{Mo}$ undergoes radioactive decay by emitting a single beta particle. Which of the following reactions corresponds to this process?

1. $^{99}_{42}\text{Mo} \longrightarrow ^{99}_{43}\text{Tc} + ^0_{-1}\beta$
2. $^{99}_{42}\text{Mo} + ^0_{-1}\beta \longrightarrow ^{99}_{41}\text{Nb}$
3. $^{99}_{42}\text{Mo} + ^0_{-1}\beta \longrightarrow ^{99}_{43}\text{Tc}$

**036 10.0 points**

Consider the following generic gas phase reaction.



The value of K_p for this reaction is 107. Calculate the equilibrium partial pressure of gas Y_2 if the equilibrium partial pressures of XY_3 is 0.50 atm and X_2 is 0.15 atm?

1. 5.6 atm
2. 0.33 atm
3. 0.016 atm
4. 0.42 atm
5. 0.031 atm
6. 0.25 atm
7. 0.18 atm


037 10.0 points

Isocarboxazid (MW = 231.25 g/mol) is an organic monoamine oxidase inhibitor used to treat depression disorders. 38.00 grams of isocarboxazid are added to water to make a 350 mL aqueous solution. What is the osmotic pressure exerted by this solution across a semi-permeable membrane at 37°C?


1. 12.10 atm
2. 329.6 atm
3. 23.90 atm
4. 1.43 atm
5. 144.4 atm
6. 11.95 atm

038 10.0 points


The chlorination of methane is an exothermic reaction with a two-step mechanism shown below:




Which of the following reaction coordinate diagrams best fits this data?

**039 10.0 points**

Rank the following acids in increasing order of acidity.



1. $\text{HCN} < \text{NH}_3\text{OH}^+ < \text{HNO}_2 < \text{HBrO}$
2. $\text{NH}_3\text{OH}^+ < \text{HBrO} < \text{HCN} < \text{HNO}_2$



040 10.0 points

Calculate the pH of a 0.018 M $\text{Ba}(\text{OH})_2$ solution.

1. 5.26

2. 8.44

3. 12.56

4. 12.26

5. 1.44

6. 1.74

041 10.0 points

How does a catalyst affect the rate of a chemical reaction?

1. A catalyst increases the rate constant by increasing the activation energy

2. A catalyst increases the energy of the transition state such that a larger number of particles have sufficient energy to overcome the activation energy

3. A catalyst increases the rate constant by providing an alternate mechanism with a lower activation energy

4. A catalyst decreases the rate constant by lowering the activation energy

042 10.0 points

When direct heat is applied to potassium chlorate, KClO_3 , it decomposes to form KCl and other byproducts. Was chlorine oxidized or reduced? How many electrons were transferred during the process?

1. oxidized, 3 electrons

2. reduced, 9 electrons

3. oxidized, 6 electrons


4. oxidized, 4 electrons

5. reduced, 3 electrons

6. reduced, 6 electrons

043 10.0 points

Consider the following cell:



What is ΔG° for the overall cell reaction that is represented here? Balance the reaction using the lowest possible integer values.

1. -91.2 kJ

2. $+91.2$ kJ

3. $+877$ kJ

4. $+182$ kJ

5. -877 kJ

6. -182 kJ

044 10.0 points

What is the mass of the barium chromate precipitate resulting from the addition of 300 mL 0.025 M $\text{Ba}(\text{OH})_2$ to 200 mL 0.040 M Na_2CrO_4 ?

1. 1.90 g

2. 3.80 g


3. 2.03 g

4. 2.52 g

5. 3.48 g

045 (part 1 of 3) 10.0 points

A sample of 30 mL of a weak acid (HA) solution was titrated with 0.075 M NaOH. The pH curve for this titration is shown.



What is the concentration of the original weak acid solution (the 30 mL) ?

1. 0.075 M
2. 0.055 M
3. 0.032 M
4. 0.048 M
5. 0.022 M

046 (part 2 of 3) 10.0 points

Which of the following is the value of K_a for the weak acid, HA ?

1. 1.2×10^{-7}
2. 5.0×10^{-7}
3. 1.3×10^{-6}
4. 7.6×10^{-5}
5. 3.2×10^{-10}

047 (part 3 of 3) 10.0 points

Below is a listing of five indicators and their associated pK_a values. Which indicator would be the best one to use for this titration?

1. bromocresol purple, 6.4
2. methyl red, 5.0
3. thymol blue, 9.3
4. bromophenol blue, 4.1
5. phenol red, 7.4
6. alizarin yellow, 10.9

048 10.0 points

You mix 0.02 mmol $\text{Sr}(\text{NO}_3)_2$ solution and 0.05 mmol NaF solution to form a 100 mL solution. What precipitate (if any) forms?

1. NaNO_3
2. $\text{Sr}(\text{NO}_3)_2$
3. NaF
4. SrF_2
5. No precipitate forms

049 10.0 points

The enthalpy of vaporization of a liquid is measured to be about 28.4 kJ/mol and its normal boiling point is 128°C . At what temperature is the partial pressure of this substance 1180 torr?

1. 176°C
2. 150°C
3. -281°C
4. 181°C
5. 162°C
6. -381°C
7. -7.92°C

050 10.0 points

Consider a standard voltaic cell at equilibrium. Which of the following is true?

1. $E > 0$, $\Delta G > 0$, $K > 1$
2. $E < 0$, $\Delta G > 0$, $K < 0$
3. $E < 0$, $\Delta G > 0$, $K < 1$
4. $E = 0$, $\Delta G = 0$, $K = 1$
5. $E = 0$, $\Delta G = 0$, $K > 1$