version: 103 Extra Practice 1 - S20 - McCord - ch305

1	_																18
1																	2
H 1.008	2											13	14	15	16	17	He 4.003
3	4	l										5	6	7	8	9	10
Li	Ве											[°] В	С	N	o	F	Ne
6.941	9.012											10.81	12.01	14.01	16.00	19.00	20.18
11	12											13	14	15	16	17	18
Na	Mg											Al	Si	Р	S	CI	Ar
22.99	24.31	3	4	5	6	7	8	9	10	11	12	26.98	28.09	30.97	32.07	35.45	39.95
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
39.10	40.08	44.96	47.87	50.94	52.00	54.94	55.85	58.93	58.69	63.55	65.38	69.72	72.64	74.92	78.96	79.90	83.80
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rb	Sr	Y	Zr	Nb	Мо	Тс	Ru	Rh	Pd	Ag	Cd	ln	Sn	Sb	Те		Xe
85.47	87.62	88.91	91.22	92.91	95.94	(98)	101.07	102.91	106.42	107.87	112.41	114.82	118.71	121.76	127.60	126.90	131.29
55	56	57	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs	Ва	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	TI	Pb	Bi	Po	At	Rn
132.91	137.33	138.91	178.49	180.95	183.84	186.21	190.23	192.22	195.08	196.97	200.59	204.38	207.20	208.98	(209)	(210)	(222)
87_	88	89	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118
Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Cn	Nh	FI	Mc	Lv	Ts	Og
(223)	(226)	(227)	(267)	(268)	(269)	(270)	(270)	(278)	(281)	(282)	(285)	(286)	(289)	(290)	(293)	(294)	(294)

58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu
140.12	140.91	144.24	(145)	150.36	151.96	157.25	158.93	162.50	164.93	167.26	168.93	173.04	174.97
90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
232.04	231.04	238.03	(237)	(244)	(243)	(247)	(247)	(251)	(252)	(257)	(258)	(259)	(266)

constants

 $R=0.08206~\mathrm{L~atm/mol~K}$

R = 8.314 J/mol K

 $N_{\rm A} = 6.022 \times 10^{23} \ / {\rm mol}$

 $h=6.626\times 10^{-34}~\mathrm{J\cdot s}$

 $c = 3.00 \times 10^8 \text{ m/s}$

 $g = 9.81 \text{ m/s}^2$

conversions

1 atm = 760 torr

1 atm = 101325 Pa

1 atm = 1.01325 bar

 $1 \text{ bar} = 10^5 \text{ Pa}$

 $^{\circ}F = ^{\circ}C(1.8) + 32$

 $K = {}^{\circ}C + 273.15$

conversions

1 in = 2.54 cm

1 ft = 12 in

1 yd = 3 ft

1 mi = 5280 ft

1 lb = 453.6 g

1 ton = 2000 lbs

1 tonne = 1000 kg

1 gal = 3.785 L

1 gai — 0.100 L

 $1 \text{ gal} = 231 \text{ in}^3$ 1 gal = 128 fl oz

1 fl oz = 29.57 mL

water data

 $C_{\rm s,ice} = 2.09 \text{ J/g }^{\circ}\text{C}$

 $C_{\rm s,water} = 4.184 \text{ J/g} \,^{\circ}\text{C}$

 $C_{\mathrm{s,steam}} = 2.03 \mathrm{\ J/g\ ^{\circ}C}$

 $\rho_{\rm water} = 1.00 \text{ g/mL}$

 $\rho_{\rm ice} = 0.9167 \ {\rm g/mL}$

 $\rho_{\rm seawater} = 1.024 \text{ g/mL}$

 $\Delta H_{\rm fus} = 334 \text{ J/g}$

 $\Delta H_{\rm vap} = 2260 \text{ J/g}$

 $K_{\rm w} = 1.0 \times 10^{-14}$

This extra practice set can be used to test your knowledge for the upcoming exam.

- 1. Most weak acids and weak bases fall into which range of ionization given below when they are dissolved into water?
- a. 1% or less
- b. just below 100%
- c. 60% to 75%
- d. 30% to 50%
- e. 10% to 20%
- 2. A titration is performed by adding a strong acid to fully neutralize a weak base analyte. The titration is run exactly to the equivalence point (where there are equal moles of the strong acid added and the initial weak base solution). What is the generic reaction for this experiment?
- a. Strong Acid + Weak Base \longrightarrow Weak Base + Water
- b. Strong Acid + Weak Base \longrightarrow Strong Acid + Water
- c. Strong Acid + Weak Base \longrightarrow Strong Base + Water
- d. Strong Acid + Weak Base \longrightarrow Weak Acid + Water
- **3.** When writing out the expression for an equilibrium constant, which of the species listed are left *out* of the expression?
- a. aqueous species
- b. solids
- c. gases
- **4.** Solution A has a hydronium ion concentration, $[H_3O^+]$, that is one thousand times higher than Solution B at identical temperature and pressure. Which of the following statements must be true?
- a. Solution A has a pH that is 3 units greater than Solution B.
- b. Solution A can have the same pH as Solution B.
- c. Solution A has a pH that is 1/1000 times that of Solution B.
- d. Solution A has a pH that is 3 units less than Solution B.

- **5**. What is $[OH^-]$ in a 0.025 M solution of $Sr(OH)_2$?
- a. 0.013 M
- b. 0.025 M
- c. 0.050 M
- d. approximately 1% or less of 0.025 M
- 6. Calculate the pH of a 0.0337 M HNO₃ solution.
- a. 1.472
- b. 5.599
- c. 0.925
- d. 8.529
- e. 0.0337
- 7. A 0.020 M solution of citric acid has a pH of 2.46. A 0.020 M solution of ascorbic acid has a pH of 2.93. Which of these is the stronger acid?
- a. citric acid
- b. ascorbic acid
- c. You cannot tell from the information provided.
- 8. When a person hyperventilates, the primary problem is not that they are inhaling too much oxygen. Rather, they are exhaling too much carbon dioxide, which can affect blood pH. The process of carbon dioxide dissolving in blood and dissociating as carbonic acid is shown in the steps below:

$$\begin{aligned} CO_2(g) &\rightleftharpoons CO_2(aq) \\ H_2O(\ell) + CO_2(aq) &\rightleftharpoons H_2CO_3(aq) \\ H_2CO_3(aq) &\rightleftharpoons H^+(aq) + HCO_3^-(aq) \end{aligned}$$

Select the answer that best describes the effect of exhaling too much carbon dioxide.

- a. blood pH increases
- b. blood pH decreases

- **9.** A salon wants to neutralize 55.0 gallons of water that have become contaminated with facial peel acids, resulting in a pH of 3.25. What volume of 0.015 M NaOH solution are needed to neutralize the acidic waste?
- a. 1.6×10^{-8} gal
- b. 0.0375 gal
- c. 2.06 gal
- d. 3.09 gal
- e. $1 \times 10^{-3.25}$ gallons
- 10. What is the $[OH^-]$ of an aqueous solution if $[H_3O^+] = 3.31 \times 10^{-4} \text{ M}$?
- a. $6.12 \times 10^{10} \text{ M}$
- b. $3.02 \times 10^{-11} \text{ M}$
- c. $3.02 \times 10^{10} \text{ M}$
- d. $3.31 \times 10^{-18} \text{ M}$
- e. $3.31 \times 10^{-4} \text{ M}$
- 11. Identify the conjugate base of formic acid, HCOOH?
- a. COO^{2-}
- b. H_3O^+
- c. HCOO-
- d. OH-
- e. H_2O
- 12. What is the pH of a weak base solution that has a percent ionization of 1.8% at 0.25 M?
- a. 9.45
- b. 11.65
- c. 12.68
- d. 2.35
- e. 13.40

- 13. What does it mean to say that a weak acid (HA_1) has a greater K_a than a second weak acid (HA_2) ?
- a. The ratio of products to reactants for ${\rm HA}_1$ is less than it is for ${\rm HA}_2$
- b. The ratio of products to reactants for HA_1 is greater than it is for HA_2
- c. HA_1 will produce a more basic solution
- d. HA₂ must have a greater percent ionization
- 14. How does water dissolve an ionic compound?
- a. The nonpolar nature of water allows the water molecules to surround a full ionic compound without separating the charged ions.
- b. The polarity of water allows the negative pole to surround cations and the positive pole to surround anions.
- c. The nonpolar nature of water allows the water molecules to dissociate ionic compounds and nonselectively surround anions and cations.
- d. The polarity of water allows the negative pole to surround anions and the positive pole to surround cations.
- 15. Calculate the molarity of [H⁺] when 12.0 grams of HCl are dissolved to make a 1.36 gallon solution.
- a. 0.0639 M
- b. 8.82 M
- c. 15.6 M
- d. 0.013 M
- e. 0.287 M
- 16. Water is amphiprotic, which means it can act as an acid and a base. Consider the following reaction:

$$\mathrm{HCl}(\mathrm{aq}) + \mathrm{H_2O}(\ell) \rightleftharpoons \mathrm{H_3O^+}(\mathrm{aq}) + \mathrm{Cl^-}(\mathrm{aq})$$

Is water acting as an acid or a base in this reaction?

- a. acid
- b. neither
- c. base

- 17. When pure water is carbonated with CO₂ gas the pH tends to rise slightly above 7 due to the alkaline nature of carbonate.
- a. false
- b. true
- 18. Which of the following species can get into into the air and cause "acid rain"?
- a. methane
- b. ozone
- c. sulfur oxides
- d. carbon dioxide
- e. nitrogen
- 19. Which of the following species contributes most to the natural acidity of rain?
- a. nitrogen
- b. sulfur oxides
- c. carbon dioxide
- d. ozone
- e. methane
- 20. Assume that each of the following bases are mixed at the same concentration of 0.05 mol/L. Which one will give the most basic solution?

ammonia, $K_{\rm b} = 1.8 \times 10^{-5}$

hydrazine, $K_{\rm b} = 1.7 \times 10^{-6}$

methylamine, $K_{\rm b} = 1.7 \times 10^{-9}$

ethylamine, $K_{\rm b} = 5.6 \times 10^{-4}$

- a. methylamine
- b. ethylamine
- c. hydrazine
- d. All solutions have the same concentration and will therefore have the same basicity.
- e. ammonia

- 21. Select the set of compounds which contains NO strong acids.
- a. Ca(OH)₂, HI, CH₃COOH
- b. $Ca(OH)_2$, HF, CH_3COOH
- c. CH₃COOH, HNO₃, CaCO₃
- d. CaCO₃, NaCl, HCl
- e. HBr, HClO₄, CaCO₃
- 22. A weak acid, HA, has a 2.5% ionization in a 0.10 M solution. What is the pH of this solution?
- a. 0.0025
- b. 1.00
- c. 3.38
- d. 5.24
- e. 2.12
- f. 2.60
- **23.** The K_a for nitrous acid (HNO₂) is 4.0×10^{-4} . What is the K_b for nitrite, NO₂⁻?
- a. 5.2×10^{-10}
- b. 4.0×10^{-18}
- c. 2.5×10^3
- d. 2.5×10^{-11}
- e. 2.0×10^{-8}
- 24. Which of the following pH values is slightly acidic but still capable of sustaining the majority of aquatic life?
- a. pH = 7.0
- b. pH = 6.6
- c. pH = 8.6
- d. pH = 11.9
- e. pH = 2.7

25. Calculate the pOH of $0.012~\mathrm{M~Ba(OH)_2}$.

- a. 12.1
- b. 1.92
- c. 2.34
- d. 12.4
- e. 1.62

Remember to bubble in ALL your answers BEFORE time is called. Double check your name, uteid, and version number before you turn in your bubblesheet. You must keep your exam for future reference. Please do not lose it. We will not replace it.