



# HW06 - Buffers, Titrations, and Polyprotics

## Question 1 1.0 pts

When an acid and base neutralize each other, the products are generally water and...

- a. a gel.
- b. a colloid.
- c. an ion.
- d. a salt.

## Question 2 1.0 pts

How many moles of  $\text{Ca(OH)}_2$  are needed to neutralize three moles of HCl?

- a. 1.5
- b. 3
- c. 2
- d. 1


## Question 3 1.0 pts

An aqueous solution is prepared with 2 moles of HCl and 1 mole of  $\text{Ca(OH)}_2$ . The resulting solution contains mainly...

- a. water,  $\text{Cl}^-$  ions, and  $\text{Ca}^{2+}$  ions.
- b. water,  $\text{Cl}^-$  ions,  $\text{H}^+$  ions,  $\text{OH}^-$  ions, and  $\text{Ca}^{2+}$  ions.
- c. water,  $\text{Cl}^-$  ions,  $\text{OH}^-$  ions, and  $\text{Ca}^{2+}$  ions.
- d. water,  $\text{Cl}^-$  ions,  $\text{H}^+$  ions, and  $\text{Ca}^{2+}$  ions.

## Question 4 1.0 pts


Identify the products of the following chemical reaction:



- a.  $3\text{H}^+ + 3\text{O}_2 + \text{H}_3\text{Li}_3$
- b.  $\text{Li}_3\text{P} + 2\text{H}_2\text{O} + \text{H}_3\text{O}_5$
- c.  $3\text{LiH} + (\text{OH})_3\text{PO}_4$
- d.  $\text{Li}_3\text{PO}_4 + 3\text{H}_2\text{O}$

## Question 5 1.0 pts

Identify the products of the following chemical reaction:



- a.  $\text{SrH}_2 + \text{HNO}_5$
- b.  $\text{SrNO}_3 + \text{H}_2\text{O}$
- c.  $\text{Sr(NO}_2)_2 + 2\text{H}_2\text{O}_2$
- d.  $\text{Sr(NO}_3)_2 + 2\text{H}_2\text{O}$

## Question 6 1.0 pts

Aqueous ammonia can be used to neutralize sulfuric acid and nitric acid to produce two salts extensively used as fertilizers. They are...

- a. cyanamide and cellulose nitrate, respectively
- b.  $\text{NH}_4\text{SO}_4$  and  $\text{NH}_4\text{NO}_3$ , respectively
- c.  $\text{NH}_4\text{SO}_3$  and  $\text{NH}_4\text{OH}$ , respectively
- d.  $(\text{NH}_4)_2\text{SO}_4$  and  $\text{NH}_4\text{NO}_3$ , respectively

## Question 7 1.0 pts

Identify the salt that is produced from the acid-base neutralization reaction between potassium hydroxide and acetic acid.

- a. potassium cyanide
- b. potassium amide
- c. potassium formate
- d. potassium acetate

## Question 8 1.0 pts

What is the pH of an aqueous solution that is 0.018 M  $\text{C}_6\text{H}_5\text{NH}_2$  ( $K_b = 4.3 \times 10^{-10}$ ) and 0.12 M  $\text{C}_6\text{H}_5\text{NH}_3\text{Cl}$ ?

- a. 3.81
- b. 4.02
- c. 4.63
- d. 2.87

## Question 9 1.0 pts

A buffer solution is made by dissolving 0.45 moles of a weak acid (HA) and 0.33 moles of KOH into 710 mL of solution. What is the pH of this buffer?  $K_a = 6 \times 10^{-6}$  for HA.

- a. 5.22
- b. 8.34
- c. 13.23
- d. 5.66

## Question 10 1.0 pts

Which one of the following combinations is NOT a buffer solution?

- a. HBr and KBr
- b. HCN and NaCN
- c.  $\text{CH}_3\text{COOH}$  and  $\text{NaCH}_3\text{COO}$
- d.  $\text{NH}_3$  and  $(\text{NH}_4)_2\text{SO}_4$

## Question 11

1.0 pts

Which of the following mixtures will be a buffer when dissolved in a liter of water?

- 0.2 mol HBr and 0.1 mol NaOH
- 0.3 mol NaCl and 0.3 mol HCl
- 0.2 mol HF and 0.1 mol NaOH
- 0.1 mol  $\text{Ca}(\text{OH})_2$  and 0.3 mol HI

## Question 12

1.0 pts

What is the pH of a solution which is 0.600 M in dimethylamine ( $(\text{CH}_3)_2\text{NH}$ ) and 0.400 M in dimethylamine hydrochloride ( $(\text{CH}_3)_2\text{NH}_2\text{Cl}$ )?  $K_b$  for dimethylamine =  $7.4 \times 10^{-4}$ .

- 11.21
- 10.78
- 11.05
- 10.87

## Question 13

1.0 pts

What would be the final pH if 0.0100 moles of solid NaOH were added to 100 mL of a buffer solution containing 0.600 molar formic acid (ionization constant =  $1.8 \times 10^{-4}$ ) and 0.300 M sodium formate?

- 3.65
- 3.44
- 4.05
- 3.84

## Question 14

1.0 pts

A buffer was prepared by mixing 0.200 moles of ammonia ( $K_b = 1.8 \times 10^{-5}$ ) and 0.200 moles of ammonium chloride to form an aqueous solution with a total volume of 500 mL. 250 mL of the buffer was added to 50.0 mL of 1.00 M HCl. What is the pH of this second solution?

- 8.38
- 8.53
- 8.18
- 8.78

## Question 15

1.0 pts

A solution is 0.30 M in  $\text{NH}_3$ . What concentration of  $\text{NH}_4\text{Cl}$  would be required to achieve a buffer solution with a final pH of 9.0?  $K_b = 1.8 \times 10^{-5}$  for  $\text{NH}_3$ .

- 0.32 M
- 0.54 M
- 0.10 M
- 0.45 M

## Question 16

1.0 pts

What is the pH at the half-stoichiometric point for the titration of 0.22 M  $\text{HNO}_2(\text{aq})$  with 0.1 M  $\text{KOH}(\text{aq})$ ? For  $\text{HNO}_2$ ,  $K_a = 4.3 \times 10^{-4}$ .

- 2.31
- 7.00
- 2.01
- 3.37

## Question 17

1.0 pts

For the titration of 50.0 mL of 0.020 M aqueous salicylic acid with 0.020 M  $\text{KOH}(\text{aq})$ , calculate the pH after the addition of 55.0 mL of the base. For salicylic acid,  $\text{p}K_a = 2.97$ .

- 10.98
- 11.26
- 7.00
- 11.02

## Question 18

1.0 pts

Consider the titration of 50.0 mL of 0.0200 M  $\text{HClO}(\text{aq})$  with 0.100 M  $\text{NaOH}(\text{aq})$ . What is the formula of the main species in the solution after the addition of 10.0 mL of base?

- HClO
- $\text{ClO}^-$
- $\text{ClO}_2$
- NaOH

## Question 19

1.0 pts

50.0 mL of 0.0018 M aniline (a weak base) is titrated with 0.0048 M  $\text{HNO}_3$ . How many mL of the acid are required to reach the equivalence point?

- 133 mL
- This is a bad titration as  $\text{HNO}_3$  is not a strong acid.
- 18.8 mL
- 4.21 mL

## Question 20

1.0 pts

When we titrate a weak base with a strong acid, the pH at the equivalence point will be...

- It is impossible to know unless we are given the  $K_b$  of the weak base.
- pH = 0
- pH > 7
- pH < 7

## Question 21

1.0 pts


What is the pH at the equivalence point in the titration of 10.0 mL of 0.35 M unknown acid HZ with 0.200 M NaOH?  $K_a = 2.4 \times 10^{-7}$  for the unknown acid HZ

- 9.86
- 10.1
- 7.00
- 4.14

## Question 22

1.0 pts

What is the pH at the equivalence point of the titration pictures below?



- 7.0
- 9.2
- 3.4
- 1.8
- 5.5

## Question 23

1.0 pts

Look at the titration diagram in the question above. What type of titration is occurring?

- a strong base titrated with a weak acid
- a weak base titrated with a strong acid
- a strong base titrated with a strong acid
- a weak base titrated with a weak acid

## Question 24

1.0 pts

The acid form of an indicator is yellow and its anion is blue. The  $K_a$  of this indicator is  $10^{-5}$ . What will be the approximate pH range over which this indicator changes color?

- $5 < \text{pH} < 7$
- $6 < \text{pH} < 8$
- $4 < \text{pH} < 6$
- $3 < \text{pH} < 5$

## Question 25

1.0 pts

The unionized form of an acid indicator is yellow and its anion is blue. The  $K_a$  of this indicator is  $10^{-5}$ . What will be the color of the indicator in a solution of pH 3?

- orange
- blue
- yellow
- green

## Question 26

2.0 pts

Aspartic acid is a polypeptide side chain found in proteins. The  $pK_a$  of aspartic acid is 3.86. If this polypeptide were in an aqueous solution with a pH of 7, the side chain would have what charge?

- there is no way to know
- negative
- positive
- neutral

## Question 27

1.0 pts

Blood contains a buffer of carbonic acid ( $\text{H}_2\text{CO}_3$ ) and hydrogen carbonate ion ( $\text{HCO}_3^-$ ) that keeps the pH at a relatively stable 7.40. What is the ratio of  $[\text{HCO}_3^-] / [\text{H}_2\text{CO}_3]$  in blood?  $K_{a1} = 4.30 \times 10^{-7}$  for  $\text{H}_2\text{CO}_3$ . (Hint: Assume  $[\text{CO}_3^{2-}] = 0$ )

- $1.71 \times 10^{-14}$
- $3.98 \times 10^{-8}$
- 10.8
- 0.0926

## Question 28

2.0 pts

$\text{H}_2\text{SO}_4$  is a strong acid because the first proton ionizes 100%. The  $K_a$  of the second proton is  $1.1 \times 10^{-2}$ . What would be the pH of a solution that is 0.100 M  $\text{H}_2\text{SO}_4$ ? Account for the ionization of both protons.

- 0.963
- 0.955
- 2.05
- 1.00