

HW05 - Kinetics

1 0.75 points

Consider the reaction:




What is the overall order of the reaction and the order with respect to $[\text{O}_3]$?

- 3 and 2
- 2 and 2
- 1 and 2
- 1 and 3

2 0.75 points

When the reaction below:




is proceeding under conditions such that 0.015 mol/L of N_2O is being formed each second, the rate of the overall reaction is _____ and the rate of change for NO is _____.

- 0.015 M/s; 0.045 M/s
- 0.015 M/s; -0.045 M/s
- 0.030 M/s; -0.005 M/s
- 0.015 M/s; -0.005 M/s

3 0.75 points

What is the rate law for the reaction below:




if the following data were collected?

Exp	[A]	[B]	[C]	Initial Rate
1	0.4	1.2	0.7	2.32×10^{-3}
2	1.3	1.2	0.9	7.54×10^{-3}
3	0.4	4.1	0.8	9.25×10^{-2}
4	1.3	1.2	0.2	7.54×10^{-3}

- rate = $4.48 \times 10^{-3} [\text{A}] [\text{B}]^2 [\text{C}]$
- rate = $1.79 \times 10^{-3} [\text{B}]^2 [\text{C}]$
- rate = $3.36 \times 10^{-3} [\text{A}] [\text{B}]^3$
- rate = $5.37 \times 10^{-3} [\text{A}] [\text{B}]^3$
- rate = $1.49 \times 10^{-3} [\text{B}]^3 [\text{C}]$

4 0.75 points

A chemical reaction is expressed by the balanced chemical equation:



Consider the data below:

exp	[A] ₀	[B] ₀	initial rate (M/min)
1	0.15	0.15	0.00110363
2	0.15	0.3	0.0044145
3	0.3	0.3	0.008829

Find the rate law for the reaction.

- rate = $k [\text{B}]^2$
- rate = $k [\text{A}]^2 [\text{B}]$
- rate = $k [\text{A}] [\text{B}]$
- rate = $k [\text{A}] [\text{B}]^2$

5 0.75 points

Calculate the value of the rate constant (k) for the reaction in question 4.

- 0.00110
- 0.000166
- 0.327
- 0.00736


6 0.75 points

If the initial concentrations of both A and B are 0.31 M for the reaction in questions 4 and 5, at what initial rate is C formed?

- 0.101 M/min
- 0.00974 M/min
- 0.0314 M/min
- 0.00974 M/min

7 0.75 points

We know that the rate expression for the reaction below:



at a certain temperature is rate = $[\text{NO}]^2 [\text{O}_2]$. We carry out two experiments involving this reaction at the same temperature, but in the second experiment the initial concentration of NO is doubled while the initial concentration of O_2 is halved. The initial rate in the second experiment will be how many times that of the first?

- 4
- 1
- 2
- 8

8 0.75 points

Consider the data collected for a chemical reaction between compounds A and B that is first order in A and first order in B:

rxn	[A] ₀	[B] ₀	rate (M/s)
1	0.2	0.05	0.1
2	?	0.05	0.4
3	0.4	?	0.8

From the information above for 3 experiments, determine the missing concentrations of A and B. Answers should be in the order [A] then [B].

- 0.20 M; 0.80 M
- 0.40 M; 0.20 M
- 0.80 M; 0.10 M
- 0.80 M; 0.20 M
- 0.40 M; 0.10 M


9 0.75 points

For a reaction that is zero-order overall...

- the rate constant is zero.
- the reactant concentration does not change with time.
- the activation energy is zero.
- the rate does not change during the reaction.

10 0.75 points

Consider the reaction below:



If it is 1st order in A and 0th order in B, a plot of $\ln[\text{A}]$ vs time will have a slope that is...

- slowly increasing.
- decreasing exponentially.
- increasing exponentially.
- constant.

11 0.75 points

At a certain fixed temperature, the reaction below:




is found to be first order in the concentration of A and zeroth order in the concentration of B. The reaction rate constant is 0.05s^{-1} . If 2.00 moles of A and 4.00 moles of B are placed in a 1.00 liter container, how many seconds will elapse before the concentration of A has fallen to 0.30 moles/liter?

- There is not enough information to answer.
- 37.94 sec
- 2.83 sec
- 10.22 sec

12 0.75 points

The reaction below:



is observed to obey first-order kinetics. Which of the following plots should give a straight line?

- $[\text{A}]$ vs t^{-1}
- $\ln[\text{A}]$ vs k
- $\ln[\text{A}]$ vs k^{-1}
- $[\text{A}]$ vs k
- $[\text{A}]$ vs t
- $\ln[\text{A}]$ vs t^{-1}
- $\ln[\text{A}]$ vs t

13 0.75 points

A reaction is found to be first order with respect to one of the reactant species, A. When might a plot of $\ln[\text{A}]$ vs time NOT yield a straight line?

- if the reaction has any significant backward rate
- All of the other answers could be correct.
- when the rate also depends on the concentration of another reactant as well
- if the reaction comes to equilibrium

14 0.75 points

The reaction rate constant is determined to be $0.012\text{M}^{-1}\text{s}^{-1}$. If after 27 minutes the amount of A left is 0.048 M. What was the initial concentration of A?

- 19.49
- 0.049
- 2.53×10^{16}
- 0.72

15 0.75 points

For the reaction below:



at 800K, a plot of $\ln[\text{cyclobutane}]$ vs t gives a straight line with a slope of -1.6s^{-1} . Calculate the time needed for the concentration of cyclobutane to fall to 1/16 of its initial value.

- 1.7 sec
- 0.63 sec
- 1.6 sec
- 1.3 sec


16 0.75 points

The initial concentration of the reactant A in a first-order reaction is 1.2 M. After 69.3 sec, the concentration has fallen to 0.3 M. What is the rate constant k ?

- 0.01s^{-1}
- not enough information
- 0.2s^{-1}
- 0.02s^{-1}

17 0.75 points

Consider the reaction below:






If it has a half-life of 1.6 sec, how long will it take a system with $[\text{H}_2\text{CO}_3]_0$ of 2M to reach $[\text{H}_2\text{CO}_3]$ of 125mM?

- 2.9 sec
- Not enough information is given.
- 6.4 sec
- 3.2 sec

18 0.75 points

Consider the following elementary reactions:




Identify the molecularity of each reaction respectively.

- it is impossible to know without knowing the overall reaction for each
- all three elementary reactions are bimolecular
- tetramolecular, termolecular, pentamolecular
- bimolecular, unimolecular, termolecular

19 0.75 points

A and B react to form C according to the single step reaction below:




Which of the following is the correct rate equation for [B] and the correct units for the rate constant of this reaction?

- $\frac{\Delta[\text{B}]}{\Delta t} = -k[\text{A}][\text{B}]^2; \frac{1}{\text{M}^2\text{s}}$
- $\frac{\Delta[\text{B}]}{\Delta t} = -2k[\text{A}][\text{B}]; \frac{1}{\text{M}\cdot\text{s}}$
- $\frac{\Delta[\text{B}]}{\Delta t} = -2k[\text{A}][\text{B}]^2; \frac{1}{\text{M}^2\cdot\text{s}}$
- $\frac{\Delta[\text{B}]}{\Delta t} = -\frac{2k[\text{A}][\text{B}]}{[\text{C}]}; \frac{1}{\text{M}\cdot\text{s}}$

20 0.75 points

Consider the mechanism below:



What is the rate law?

- rate = $k_1[\text{NO}_2][\text{F}_2]$
- rate = $k_2[\text{NO}_2][\text{F}]$
- rate = $k_1k_2[\text{NO}_2]^2$
- rate = $k_1[\text{NO}_2\text{F}][\text{F}_2]$
- rate = $k_2[\text{NO}_2]^2$

21 1 point

Determine the overall balanced equation for a reaction having the following proposed mechanism:

and write an acceptable rate law.

- $\text{A} + \text{B}_2 \rightarrow \text{C}_2 + \text{D}; \text{rate} = k[\text{A}][\text{B}_2]$
- $\text{E}_3 + \text{A} \rightarrow \text{B}_2 + \text{C}_2; \text{rate} = k[\text{E}_3][\text{A}]$
- $\text{A} + 2\text{B}_2 \rightarrow \text{C}_2 + \text{D}; \text{rate} = k[\text{B}_2]^2$
- $2\text{B}_2 \rightarrow \text{E}_3 + \text{D}; \text{rate} = k[\text{B}_2]^2$

22 1 point

Consider the reaction below:

The proposed mechanism of this reaction is:

What is the rate of the overall reaction?

- rate = $\frac{k_1k_2}{k_{-1}}[\text{I}_2][\text{H}_2]$
- rate = $\frac{k_1k_2}{k_{-1}}[\text{I}_2][\text{H}_2]$
- rate = $\frac{k_1k_2}{k_{-1}}[\text{I}]^2[\text{H}_2]$
- rate = $k_1k_2[\text{I}_2][\text{H}_2]$
- rate = $k_2[\text{I}]^2[\text{H}_2]$

23 1 point

A reaction rate increases by a factor of 655 in the presence of a catalyst at 37°C. The activation energy of the original pathway is 106 kJ/mol. What is the activation energy of the new pathway, all other factors being equal?

- 16,600 kJ/mol
- 89.3 kJ/mol
- 89.3 J/mol
- 16,600 J/mol

24 1 point

A given reaction has an activation energy of 24.52 kJ/mol. At 25°C, the half-life is 4 minutes. At what temperature will the half-life be reduced to 20 seconds?

- 100°C
- 125°C
- 150°C
- 115°C

25 1 point

For the reaction below:

a plot of $\ln k$ vs $1/T$ gives a straight line with a slope equal to $-5.1 \times 10^3\text{K}$

What is the activation energy for this reaction?

- 42 kJ/mol
- 98 kJ/mol
- 5.1 kJ/mol
- 12 kJ/mol

26 1 point

A certain reaction has an activation energy of 0.8314 kJ/mol and a rate constant of 2.718s^{-1} at -73°C. At -173°C, which expression for the rate constant is correct?

- $\ln(k_2) = -0.5$
- $\ln(k_2) = 1.5$
- $\ln(k_2) = 1$
- $\ln(k_2) = 0.5$

27 1 point

A food substance kept at 0°C becomes rotten (as determined by a good microorganism) in 8.3 days. The same food rots in 10.6 hours at 30°C. Assuming the kinetics of the microorganism enzymatic action is responsible for the rate of decay, what is the activation energy for the decomposition process? Hint: Rate varies INVERSELY with time; a faster rate produces a shorter decomposition time.

- 0.45 kJ/mol
- 67.2 kJ/mol
- 2.34 kJ/mol
- 23.4 kJ/mol

28 1 point

A catalyst...

- changes the reaction mechanism to ensure that K is increased.
- increases K to favor product formation.
- speeds up the reaction but does not change K.
- speeds up the reaction and increases K to favor product formation.

29 1 point

All else being equal, a reaction with a higher activation energy compared to one with a lower activation energy will...

- be more endothermic.
- proceed slower.
- be more exothermic.
- proceed faster.

30 1 point

Consider the potential energy diagram below:

What is the change in enthalpy (ΔH) for the reaction $\text{A} \rightarrow \text{B}$?

- 350 kJ
- 100 kJ
- 100 kJ
- 350 kJ

31 1 point

Consider the potential energy diagram in question 13. What is the activation energy (E_a) for the reaction?

- 100 kJ
- 200 kJ
- 250 kJ
- 350 kJ

32 1 point

Which of the following statements is TRUE?

- If the exponents in the rate-law do not match the coefficients in the balanced equation, then we know that the reaction does not take place in one step.
- The rate-law for a reaction can be predicted from the balanced chemical equation.
- The exponents in the rate-law must match the coefficients in the balanced chemical equation for the reaction.
- If the exponents in the rate-law do not match the coefficients in the balanced chemical equation, then we know that the reaction takes place in one step.

33 1 point

"Reaction mechanisms usually involve only unimolecular or bimolecular steps."

Is this statement true or false?

- True, because steps of higher molecularity would not be compatible with observed reaction rate laws.
- False.
- True, because collisions of higher molecularity would occur too infrequently to account for an observed rate.
- True, because the activation