

## HW06 - Solubility

I put a couple of kinetics problems at the start of this HW for more practice on those. Note... to speed up solution entry and aid in simpler files. Scientific notation numbers will often be given in the "E" format - which is really a computer science thing. But this IS a science course, for science majors. So know that a "E" or "e" in a number stands for "times 10 to the \_\_\_ power". You could see this format anywhere, but I try to only use it in the explanations or solutions. Examples:  $6.022 \times 10^{23}$  is simply 6.022E23 or 6.022e23 or for little numbers,  $K_a = 1.8 \times 10^{-5}$  is also 1.8E-5.

1 2 points

Consider the following simple generic decomposition reaction:



Initially, 0.0800 moles of A are dissolved into 400 mL of water. There is no B initially, like most reactions. The rate constant for this reaction is  $0.0250 \text{ M}^{-1}\text{s}^{-1}$ . How long will it take to make 0.144 moles of B?

- 1 hr
- 45 min
- 10 min
- 1.5 min
- 25 min
- 30 min

2 2 points

Reactant "A" is added to a reaction chamber such that its partial pressure is 0.25 atm. The rate constant for this second order reaction is  $0.040 \text{ atm}^{-1}\text{s}^{-1}$  in reactant A. What is the half life of this reaction under these conditions? note: yes, you can, and you should work this in partial pressures (atm) and not concentration. Look at the rate constant... it is in atm.

- 17 s
- 100 s
- 36 s
- 75 s
- 124 s

3 2 points

What is the net ionic equation for the reaction between aqueous solutions of  $\text{Na}_3\text{PO}_4$  and  $\text{CuSO}_4$ ?

Note: assume all ionic species are (aq).

- $\text{Cu}^{2+}(aq) + \text{PO}_4^{3-}(aq) \longrightarrow \text{CuPO}_4(s)$
- No reaction occurs since no precipitate is formed.
- $3\text{Cu}^{2+}(aq) + 2\text{PO}_4^{3-}(aq) \longrightarrow \text{Cu}_3(\text{PO}_4)_2(s)$
- $2\text{Na}^+(aq) + \text{SO}_4^{2-}(aq) \longrightarrow \text{Na}_2\text{SO}_4(s)$

4 2 points

If you mix equal volumes and concentrations of aqueous solutions of  $\text{Cu}(\text{NO}_3)_2$  and  $\text{K}_2\text{S}$ , which ions will remain in solution as spectators in the reaction?

- $\text{Cu}^{2+}, \text{S}^{2-}$
- $\text{K}^+, \text{NO}_3^-$
- No ions are present as both products form precipitates.
- $\text{Cu}^{2+}, \text{NO}_3^-, \text{K}^+, \text{S}^{2-}$

5 1 point

Molar solubility is...

- the total molarity of the solution.
- the number of grams of a compound that dissolve to give one liter of saturated solution.
- equal to the  $K_{sp}$ .
- the number of moles of a compound that dissolve to give one liter of saturated solution.

6 2 points

A sample of pure water is saturated with  $\text{PbCl}_2$ . In this saturated solution, which of the following is true?

- $[\text{Pb}^{2+}] = [\text{Cl}^-]$
- $K_{sp} = [\text{Pb}^{2+}]^2[\text{Cl}^-]$
- $K_{sp} = [\text{Pb}^{2+}][\text{Cl}^-]$
- $[\text{Pb}^{2+}] = 0.5[\text{Cl}^-]$

7 2 points

A hypothetical ionic substance  $\text{M}_3\text{X}_2$  dissociates in water to form  $\text{M}^{2+}$  and  $\text{X}^{3-}$  ions.

The molar solubility of  $\text{M}_3\text{X}_2$  is  $4.5 \times 10^{-3} \text{ mol/L}$ .

What is the value of the solubility product constant?

- $2.0 \times 10^{-5}$
- $2.0 \times 10^{-10}$
- $1.1 \times 10^{-8}$
- $1.8 \times 10^{-12}$
- $3.6 \times 10^{-7}$
- $8.3 \times 10^{-15}$

8 2 points

The value of  $K_{sp}$  for  $\text{SrSO}_4$  is  $2.8 \times 10^{-7}$ . What is the molar solubility of  $\text{SrSO}_4$ ?

- $1.4 \times 10^{-7} \text{ mol/L}$
- $7.6 \times 10^{-7} \text{ mol/L}$
- $5.3 \times 10^{-4} \text{ mol/L}$
- $2.8 \times 10^{-7} \text{ mol/L}$

9 2 points

Determine the molar solubility of a salt with the generic formula  $\text{AB}_2$  if  $K_{sp} = 3.7 \times 10^{-5}$ .

- 0.063 M
- 0.0061 M
- 0.021 M
- 0.033 M
- 0.052 M

10 1 point

Rank the following salts from lowest to highest molar solubility:

$\text{AgI}$	$K_{sp} = 8.5 \times 10^{-17}$
$\text{Cd}_3(\text{AsO}_4)_2$	$K_{sp} = 2.2 \times 10^{-33}$
$\text{AlPO}_4$	$K_{sp} = 9.8 \times 10^{-21}$
$\text{CaSO}_4$	$K_{sp} = 4.9 \times 10^{-5}$

- $\text{Cd}_3(\text{AsO}_4)_2 < \text{AgI} < \text{AlPO}_4 < \text{CaSO}_4$
- $\text{CaSO}_4 < \text{AgI} < \text{AlPO}_4 < \text{Cd}_3(\text{AsO}_4)_2$
- $\text{Cd}_3(\text{AsO}_4)_2 < \text{AlPO}_4 < \text{AgI} < \text{CaSO}_4$
- $\text{AlPO}_4 < \text{AgI} < \text{Cd}_3(\text{AsO}_4)_2 < \text{CaSO}_4$

11 2 points

A hypothetical compound  $\text{MX}_3$  with a molar mass of 125 g/mol has a solubility of 1.5 g/L. What is the value of  $K_{\text{sp}}$  for  $\text{MX}_3$ ?

- $1.4 \times 10^{-4}$
- $6.9 \times 10^{-6}$
- $5.6 \times 10^{-7}$
- $1.2 \times 10^{-2}$
- $3.6 \times 10^{-8}$
- $1.2 \times 10^{-9}$

12 1 point

Determine if a precipitate will form when 0.96 g  $\text{Na}_2\text{CO}_3$  is combined with 0.20 g  $\text{BaBr}_2$  in a 10 L solution. (For  $\text{BaCO}_3$ ,  $K_{\text{sp}} = 2.8 \times 10^{-9}$ ).

- $\text{BaBr}_2$  will remain in solid form as it is insoluble in water.
- $\text{BaCO}_3$  does not precipitate
- It is impossible to know if any  $\text{BaCO}_3$  will precipitate with the information given.
- $\text{BaCO}_3$  precipitates

13 1 point

$\text{CaSO}_4$  has a  $K_{\text{sp}} = 3 \times 10^{-5}$ . In which of the following would  $\text{CaSO}_4$  be the most soluble?

- 1.0 M  $\text{CaCl}_2(\text{aq})$
- 0.5 M  $\text{K}_2\text{SO}_4(\text{aq})$
- $\text{CaSO}_4$  would have the same solubility in all three of these solutions
- pure water

14 2 points

A solution of  $\text{AgI}$  contains 0.019 M  $\text{Ag}^+$ . The  $K_{\text{sp}}$  of  $\text{AgI}$  is  $8.3 \times 10^{-17}$ . What is the maximum  $\text{I}^-$  concentration that can exist in this solution before a precipitate begins to form?

- $6.8 \times 10^{-9}$  M
- 0.019 M
- $1.6 \times 10^{-18}$  M
- $4.4 \times 10^{-15}$  M
- $8.3 \times 10^{-15}$  M

15 2 points

What is the molar solubility of  $\text{Zn}(\text{OH})_2$  ( $K_{\text{sp}} = 3.0 \times 10^{-17}$ ) in pure water?

(note: you might want to compare the answer on this to the one in the next question)

- $5.3 \times 10^{-13}$  M
- $8.1 \times 10^{-15}$  M
- $7.2 \times 10^{-8}$  M
- $9.5 \times 10^{-9}$  M
- $2.0 \times 10^{-6}$  M

16 2 points

What is the molar solubility of  $\text{Zn}(\text{OH})_2$  ( $K_{\text{sp}} = 3.0 \times 10^{-17}$ ) in a solution that is buffered to a pH of 9.75?

(note: you might want to compare the answer on this to the one in the previous question)

- $7.2 \times 10^{-8}$  M
- $8.1 \times 10^{-15}$  M
- $5.3 \times 10^{-13}$  M
- $9.5 \times 10^{-9}$  M
- $2.0 \times 10^{-6}$  M

17 2 points

A relatively insoluble metal hydroxide,  $\text{M}(\text{OH})_2$ , has a  $K_{\text{sp}}$  value of  $4 \times 10^{-15}$  at a particular temperature. What is the pH of a saturated solution of  $\text{M}(\text{OH})_2$ ?

- 6.80
- 8.25
- 10.40
- 9.56
- 9.30
- 9.00
- 8.72